

A WALL-CROSSING FORMULA FOR GROMOV-WITTEN INVARIANTS UNDER VARIATION OF GIT QUOTIENT

EDUARDO GONZALEZ AND CHRIS T. WOODWARD

ABSTRACT. We prove a quantum version of Kalkman's wall-crossing formula [34], [41] comparing intersection pairings on geometric invariant theory (git) quotients related by a change in polarization, under certain stable=semistable assumptions. The formula is the same, but each expression in the formula is quantized in the sense that it is replaced by an integral over moduli spaces of certain stable maps; in particular, the wall-crossing terms are gauged Gromov-Witten invariants with smaller structure group. As an application, we show that the graph Gromov-Witten potentials of quotients related by wall-crossings of crepant type are equivalent up to a distribution in the quantum parameter that is almost everywhere zero. This is a version of the crepant transformation conjecture of Li-Ruan [43], Bryan-Graber [10], Coates-Ruan [13] etc. in cases where the crepant transformation is obtained by variation of git.

CONTENTS

1. Introduction	1
2. Kalkman's wall-crossing formula	16
3. Wall-crossing for Gromov-Witten invariants	24
4. Invariance under crepant transformations	34
References	39

1. INTRODUCTION

According to the geometric invariant theory introduced by Mumford [46], the *git quotient* $X//G$ of an action of a complex reductive group G on a projective variety X equipped with a polarization (ample equivariant line bundle) has coordinate ring equal to the G -invariant part of the coordinate ring on X . Geometrically $X//G$ is the quotient of an open *semistable locus* $X^{\text{ss}} \subset X$ by an equivalence relation, where a point $x \in X$ is semistable if there is a non-constant invariant section of a tensor power of the polarization that is non-zero at the point. If the action of G on the

Partially supported by NSF grants DMS-1104670 and DMS-0904358.

semistable locus has only finite stabilizers, then $X//G$ is the quotient of X^{ss} by the action of G , by which we mean here the *stack-theoretic* quotient, see for example [18]. If X is in addition smooth, then $X//G$ is a smooth proper Deligne-Mumford stack with projective coarse moduli space. In Kempf-Ness [36], see also Mumford et al [46], the coarse moduli space of the git quotient is identified with the symplectic quotient of X by a maximal compact subgroup of G .

The question of how the git quotient depends on the polarization, or equivalently, choice of moment map is studied in a series of papers by Guillemin-Sternberg [28], Brion-Procesi [7], Dolgachev-Hu [20], and Thaddeus [52]. Under suitable stable=semistable conditions, the git quotient $X//G$ undergoes a sequence of weighted blow-ups and blow-downs. The class of birational equivalences which appear via variation of git is reasonably large. In fact, for a class of so-called *Mori dream spaces*, any birational equivalence can be written as a composition of birational equivalences induced by variation of git [31, 45].

The question of how the cohomology of the quotient depends on the polarization is studied in Kalkman [34], which proves a wall-crossing formula for the intersection pairings under variation of symplectic quotient. Similar results, in the context of Donaldson theory, are given in Ellingsrud-Göttsche [22]. To state Kalkman's wall-crossing formula, recall that any Lie group G has a homotopically-trivial *universal bundle* $EG \rightarrow BG$, uniquely defined up to homotopy equivalence. If X is a G -space with a reasonable homotopy type then the equivariant cohomology $H_G(X)$ is the cohomology $H(EG \times_G X)$ of $EG \times_G X$. In this paper we work with rational coefficients; if G acts with only finite stabilizers then the map $EG \times_G X \rightarrow X/G$ has cohomologically trivial fibers and so $H_G(X) \cong H(X/G)$ by the Serre spectral sequence. If X is a smooth projective G -variety as above such that G acts locally freely on the semistable locus (that is, with finite stabilizers) then there is a natural map

$$\kappa_X^G : H_G(X) \rightarrow H(X//G)$$

studied in Kirwan's thesis [38], given by restriction to the semistable locus $H_G(X) \rightarrow H_G(X^{\text{ss}})$ and then descent under the quotient $H_G(X^{\text{ss}}) \cong H(X//G)$. Consider the simplest case $G = \mathbb{C}^\times$. Let $X//_\pm G$ denote the git quotients corresponding to polarizations $L_\pm \rightarrow X_\pm$. Let $\kappa_{X,\pm}^G : H_G(X) \rightarrow H(X//_\pm G)$ be the Kirwan maps and let

$$\tau_{X//_\pm G} : H(X//_\pm G) \rightarrow \mathbb{Q}, \quad \alpha \mapsto \int_{[X//_\pm G]} \alpha$$

denote integration over $X//_\pm G$. Kalkman's formula expresses the difference between the integrals $\tau_{X//_\pm G} \circ \kappa_{X,\pm}^G$ as a sum of fixed point contributions from components $X^{G,t} \subset X^G$ that are semistable for $L_t := L_-^{(1-t)/2} L_+^{(1+t)/2}$ for some $t \in (-1, 1)$. Each such fixed point has a contribution to the localization formula for the integral of a class $\alpha \in H_G(X)$ over X given as follows. Let $\nu_{X^{G,t}} \rightarrow X^{G,t}$ denote the normal bundle of the inclusion $X^{G,t} \rightarrow X$, and $\text{Eul}_G(\nu_{X^{G,t}}) \in H_G(X^{G,t})$ its G -equivariant Euler class, or equivalently, equivariant Chern class of degree equal to the real rank. We identify $H(BG)$ with the polynomial ring $\mathbb{Q}[\xi]$ in a single element ξ of degree 2, representing

the hyperplane class in the cohomology of $BG = \mathbb{C}P^\infty$. If $m_t = \text{codim}(X^{G,t})$ and

$$(1) \quad \nu_{X^{G,t}} = \bigoplus_{i=1}^{m_t} \nu_{X^{G,t},i}$$

is a decomposition into line bundles with weights $\mu_i \in \mathbb{Z}$ then

$$\text{Eul}_G(\nu_{X^{G,t}}) = \prod_{i=1}^{m_t} \text{Eul}_G(\nu_{X^{G,t},i}) = \prod_{i=1}^{m_t} (c_1(\nu_{X^{G,t},i}) + \mu_i \xi).$$

Since G acts with no non-trivial fixed vectors on $\nu_{X^{G,t}}$, the Euler class has an inverse

$$\text{Eul}_G(\nu_{X^{G,t}})^{-1} \in H(X^{G,t})[\xi, \xi^{-1}]$$

after inverting the equivariant parameter ξ . If one has a splitting as in (1) then the inverted class admits an expansion

$$\begin{aligned} \text{Eul}_G(\nu_{X^{G,t}})^{-1} &= \prod_{i=1}^{m_t} (\mu_i \xi)^{-1} (1 + c_1(\nu_{X^{G,t},i})/\mu_i \xi)^{-1} \\ &= \prod_{i=1}^{m_t} (\mu_i \xi)^{-1} (1 - c_1(\nu_{X^{G,t},i})/\mu_i \xi + (c_1(\nu_{X^{G,t},i})/\mu_i \xi)^2 - \dots). \end{aligned}$$

The contribution from a fixed point component $X^{G,t} \subset X^G$ is the integral of the restriction $\iota_{X^{G,t}}^* \alpha$ times the inverted Euler class, denoted

$$\tau_{X^{G,t}}^G : H_G(X) \rightarrow \mathbb{Q}[\xi, \xi^{-1}], \quad \alpha \mapsto \int_{[X^{G,t}]} \iota_{X^{G,t}}^* \alpha \cup \text{Eul}_G(\nu_{X^{G,t}})^{-1}.$$

Let Resid denote the residue, that is, the coefficient of ξ^{-1} :

$$\text{Resid}_\xi : \mathbb{Q}[\xi, \xi^{-1}] \rightarrow \mathbb{Q}, \quad \sum_{n \in \mathbb{Z}} a_n \xi^n \mapsto a_{-1}.$$

Theorem 1.1 (Kalkman wall-crossing formula, circle group case). *Let $G = \mathbb{C}^\times$ and let X be a smooth projective G -variety equipped with polarizations $L_\pm \rightarrow X$. Suppose that stable=semistable for both polarizations L_+, L_- . Then*

$$(2) \quad \tau_{X/+G} \kappa_{X,+}^G - \tau_{X/-G} \kappa_{X,-}^G = \sum_{t \in (-1,1)} \text{Resid}_\xi \tau_{X^{G,t}}^G.$$

In other words, failure of the following square to commute is measured by an explicit sum of wall-crossing terms:

$$\begin{array}{ccc}
& H_G(X) & \\
\swarrow \kappa_{X,+}^G & & \searrow \kappa_{X,-}^G \\
H(X//_+G) & & H(X//_-G) \\
\searrow \tau_{X/+G} & & \swarrow \tau_{X/-G} \\
& \mathbb{Q} &
\end{array}$$

The formula (2) also holds for certain quasi-projective varieties, such as vector spaces whose weights are contained in an open half-space, see the more general Theorem 2.4 below.

Example 1.2. (Integration over projective space) The following simple example illustrates the notation involved. Let $G = \mathbb{C}^\times$ act on $X = \mathbb{C}^k$ by scalar multiplication, so that $H_G(X) = \mathbb{Q}[\xi]$. Suppose that polarizations $L_\pm$ correspond to the characters ± 1 , so that the git quotients $X//_-G = \emptyset$ and $X//_+G = \mathbb{P}^{k-1}$ with a unique singular value $t = 0$ corresponding to the point $0 \in X$. The Kirwan map $\kappa_X^G : H_G(X) \rightarrow H(X//G)$ sends the generator $\xi \in H_G^2(X)$ to the hyperplane class $\omega \in H^2(X//G)$. We compute the integrals $\int_{\mathbb{P}^{k-1}} \omega^a$ for $a \in \mathbb{Z}_{\geq 0}$ via wall-crossing. On one side, the integral is zero, since the quotient is empty. By the Kalkman formula (2)

$$\begin{aligned}
\int_{\mathbb{P}^{k-1}} \omega^a &= \text{Resid}_\xi \int_{[0]} \xi^a \cup \text{Eul}_G(\mathbb{C}^k)^{-1} \\
&= \text{Resid}_\xi \xi^a / \xi^k = \begin{cases} 1 & a = k-1 \\ 0 & \text{otherwise} \end{cases}
\end{aligned}$$

confirming that ω^{k-1} is the dual of the fundamental class. This ends the example.

The main result of this paper is a generalization of Theorem 1.1 to the setting of Gromov-Witten theory, that is, quantum cohomology. The formula is the same, but the Kirwan map, trace, and fixed point contributions are *quantized* in the sense that each is replaced by a formal map depending on a formal variable q whose restriction to $q = 0$ gives the classical version above. First we recall the definition of quantum cohomology of a smooth polarized projective G -variety X . The *equivariant Novikov field* $\Lambda_X^G \subset \text{Hom}(H_2^G(X, \mathbb{Q}), \mathbb{Q})$ associated to a G -variety X with polarization $L \rightarrow X$ consists of linear combinations of formal symbols q^d for $d \in H_2^G(X, \mathbb{Q})$:

$$\Lambda_X^G := \left\{ \sum_{d \in H_2^G(X, \mathbb{Q})} c_d q^d, \quad \forall e > 0, \# \{c_d | \langle d, c_1^G(L) \rangle < e\} < \infty \right\}.$$

We write $\Lambda_{X,L}^G$ if we wish to emphasize the dependence on L . The *equivariant quantum cohomology* is the tensor product

$$QH_G(X) := H_G(X) \otimes \Lambda_X^G$$

or $QH_G(X, L)$ if we wish to emphasize the dependence on L . A more standard definition in algebraic geometry would use the cone of effective curve classes, but the definition give here has better invariance properties, for example, under Hamiltonian perturbation. Below we will need several variations on this definition. Let $\Lambda_X^{G, \text{fin}} \subset \Lambda_X^G$ denote the space of finite linear combinations of the symbols q^d . We denote by $QH_{G, \text{fin}}(X) \subset QH_G(X)$ the subspace $H_G(X) \otimes \Lambda_X^{G, \text{fin}}$. Let $\Lambda_X^{G, \geq 0} \subset \Lambda_X^G$ be the space consisting of expressions involving only powers q^d with $\langle d, c_1^G(L) \rangle \geq 0$. We denote by $QH_{G, \geq 0}(X) \subset QH_G(X)$ the subspace $H_G(X) \otimes \Lambda_X^{G, \geq 0}$.

The quantum cohomology $QH_G(X)$ has the structure of a Frobenius manifold, in particular, a family of products

$$\star_\alpha : T_\alpha QH_G(X)^2 \rightarrow T_\alpha QH_G(X), \quad \alpha \in QH_G(X)$$

defined by equivariant virtual enumeration of genus zero *stable maps*, that is, maps $u : C \rightarrow X$ from projective nodal curves C of arithmetic genus zero to X with no infinitesimal automorphisms. The moduli stack $\overline{\mathcal{M}}_{g,n}(X, d)$ of such maps of homology class $d \in H_2(X, \mathbb{Z})$ and arithmetic genus g is a proper Deligne-Mumford stack equipped with a perfect relative obstruction theory and evaluation map

$$\text{ev} = (\text{ev}_1, \dots, \text{ev}_n) : \overline{\mathcal{M}}_{g,n}(X, d) \rightarrow X^n.$$

The action of G on X induces an action of G on $\overline{\mathcal{M}}_{g,n}(X, d)$ by translation; then the evaluation maps induce a map

$$\text{ev}^* : H_G(X)^n \rightarrow H_G(\overline{\mathcal{M}}_{g,n}(X, d)).$$

The quantum product at $\alpha \in QH_G(X)$ is defined by restricting to $g = 0$ and defining for $\beta, \gamma \in H_G(X) \subset T_\alpha QH_G(X)$

$$\beta \star_\alpha \gamma := \sum_{n \geq 0} \sum_{d \in H_2^G(X, \mathbb{Z})} (q^d/n!) \text{ev}_{n+3,*} \text{ev}_1^* \alpha \cup \dots \cup \text{ev}_n^* \alpha \cup \text{ev}_{n+1}^* \beta \cup \text{ev}_{n+2}^* \gamma$$

where the push-forwards are defined using the Behrend-Fantechi virtual fundamental classes $[\overline{\mathcal{M}}_{g,n}(X, d)]_G$, and the definition is extended to $T_\alpha QH_G(X) \cong QH_G(X)$ by linearity over Λ_X^G , see [27]. Note that we take the virtual fundamental classes to lie in the homology $H_G(\overline{\mathcal{M}}_{g,n}(X, d))$ of the coarse moduli space, rather than in the Chow ring as in [27]. The quantum cohomology $QH_G(X)$ can also be defined using the smaller ring $\Lambda_X^{G, \geq 0}$ but not $\Lambda_X^{G, \text{fin}}$, because of the infinite sum.

The orbifold quantum cohomology $QH(X//G)$ of the quotient $X//G$ is defined by virtual enumeration of stable maps from orbifold curves, as follows. For any element $g \in G$ let Z_g be the centralizer of $g \in G$, $[g]$ its conjugacy class, and $\langle g \rangle \subset Z_g$ the subgroup generated by g . We denote by $\overline{I}_{X/G}$ the *rigidified inertia stack* from Abramovich-Graber-Vistoli [1], given by

$$\overline{I}_{X/G} = \bigcup_{[g]} X^{g, \text{ss}} / (Z_g / \langle g \rangle).$$

Denote by

$$QH(X//G) = H(\overline{I}_{X/G}) \otimes \Lambda_X^G$$

the quantum cohomology of the git quotient $X//G$ defined using the same Novikov field Λ_X^G ; this larger ring contains $\Lambda_{X/G}$ by virtue of Kirwan's injection $H_2(X//G) \rightarrow H_2^G(X)$. Virtual enumeration of representable morphisms from orbifold curves satisfying certain conditions to $X//G$ defines a family of products

$$\star_\alpha : T_\alpha QH(X//G)^2 \rightarrow T_\alpha QH(X//G), \quad \alpha \in QH(X//G).$$

A quantum version of the Kirwan map

$$\kappa_X^G : QH_G(X) \rightarrow QH(X//G)$$

(we keep the same notation as in the classical case) was constructed in [55]. The map κ_X^G is a formal, non-linear map with the property that each linearization

$$D_\alpha \kappa_X^G : T_\alpha QH_G(X) \rightarrow T_{\kappa_X^G(\alpha)} QH(X//G), \quad \alpha \in QH_G(X)$$

is a $\star$ -homomorphism, defined by virtual enumeration of *affine gauged maps*. Such a map is by definition a representable morphism $u : \mathbb{P}(1, r) \rightarrow X/G$ from a weighted projective line $\mathbb{P}(1, r)$, $r > 0$ to the quotient stack X/G mapping the stacky point at infinity $\mathbb{P}(r) \subset \mathbb{P}(1, r)$ to the semistable locus $X//G \subset X/G$. (More precisely, the domain is a smooth Deligne-Mumford stack of dimension one with a single stacky point with automorphism group of order r .) These are the algebro-geometric analogs of the *vortex bubbles* considered in Gaio-Salamon [23]. The compactified moduli stack $\overline{\mathcal{M}}_{n,1}^G(\mathbb{C}, X, d)$ of affine gauged maps of homology class $d \in H_2^G(X, \mathbb{Q})$ is, if stable=semistable for X , a proper smooth Deligne-Mumford stack with a perfect relative obstruction theory over the complexification of Stasheff's multiplihedron, and evaluation maps

$$\text{ev} \times \text{ev}_\infty : \overline{\mathcal{M}}_{n,1}^G(\mathbb{C}, X) \rightarrow (X/G)^n \times \overline{I}_{X/G}.$$

The quantum Kirwan map is defined for $\alpha \in H_G(X) \subset QH_G(X)$ by

$$\kappa_X^G(\alpha) := \sum_{n \geq 0} \sum_{d \in H_2^G(X, \mathbb{Q})} (q^d / n!) \text{ev}_{\infty,*} \text{ev}^*(\alpha, \dots, \alpha).$$

It is defined over the smaller equivariant Novikov ring $\Lambda_X^{G, \geq 0}$, but one obtains good surjectivity properties only using the Novikov field Λ_X^G , see [26]. The map κ_X^G is a quantization of Kirwan's in the sense that $D_0 \kappa_X^G|_{q=0}$ is the map studied in [38].

A quantization of the classical integration map over $X//G$ is given by the *graph potential* in Givental [24]

$$\tau_{X/G} : QH(X//G) \rightarrow \Lambda_X^G$$

defined by virtual enumeration of genus zero orbifold stable maps to $\mathbb{P} \times (X//G)$ of homology class $(1, d)$ for some $d \in H_2(X//G, \mathbb{Q})$. Let

$$\overline{\mathcal{M}}_n(C, X//G, d) := \overline{\mathcal{M}}_{0,n}(C \times (X//G), (1, d))$$

denote the stack of such maps of class $d \in H_2(X//G, \mathbb{Q})$, which we view as an element of $H_2^G(X, \mathbb{Q})$ via the inclusion of the semistable locus. The graph potential is defined

for $\alpha \in H(\bar{I}_{X/G}) \subset QH(X//G)$ by

$$\tau_{X/G}(\alpha) := \sum_{n \geq 0} \sum_{d \in H_2(X/G, \mathbb{Q})} (q^d/n!) \int_{[\overline{\mathcal{M}}_n(C, X/G, d)]} \text{ev}^*(\alpha, \dots, \alpha).$$

Again $\tau_{X/G}$ is defined over the equivariant Novikov ring $\Lambda_X^{G, \geq 0}$. The graph potential $\tau_{X/G}$ is related via the quantum Kirwan map to a *gauged Gromov-Witten potential* $\tau_X^G : QH_G(X) \rightarrow \Lambda_X^G$ defined by virtual enumeration of *gauged maps*, by which we mean morphisms from C to the quotient stack X/G , satisfying a Mundet stability condition [47] generalizing semistability for vector bundles on curves:

Definition 1.3. (Mundet semistability) A *gauged map* from a smooth projective curve C to the quotient stack X/G is a morphism $u : C \rightarrow X/G$, consisting of a pair $(p : P \rightarrow C, u : C \rightarrow P(X) := P \times_G X)$, see [18]. We suppose that $\mathfrak{g}_{\mathbb{C}}$ is the complexification of $\mathfrak{g}_{\mathbb{R}}$, and $\mathfrak{g}_{\mathbb{R}}$ is equipped with an inner product invariant under the action of $G_{\mathbb{R}} = \exp(\mathfrak{g}_{\mathbb{R}})$, inducing an identification $\mathfrak{g}_{\mathbb{R}} \rightarrow \mathfrak{g}_{\mathbb{R}}^{\vee}$.

- (a) (Associated graded morphism) Given a reduction of P to a parabolic subgroup $R \subset G$ given by a section $\sigma : C \rightarrow P/R$ and an antidominant, central element λ in the Lie algebra $\mathfrak{r}$ of R , there is an *associated graded* morphism given by a bundle

$$\text{Gr}(P) \rightarrow C$$

whose structure group reduces to the Levi subgroup of P and stable section

$$\text{Gr}(u) : \hat{C} \rightarrow \text{Gr}(P)(X)$$

with domain a nodal curve $\hat{C}$. The bundle $\text{Gr}(P)$ is the limit of the bundles $P_{z, \lambda}$ associated to the homomorphisms

$$R \rightarrow R, r \mapsto \text{Ad}(z^\lambda)r$$

for $z \in \mathbb{C}^\times$, while the section $\text{Gr}(u)$ is the Gromov limit of the family of sections $z^\lambda u$ of $P_{z, \lambda}(X)$ induced by λ .

- (b) (Hilbert-Mumford weight) The principal component C_0 of $\hat{C}$ is the irreducible component such that the restriction u_0 of u to C_0 maps isomorphically to C . The principal component $\text{Gr}(u)_0$ of the associated graded section $\text{Gr}(u)$ takes values in the fixed point set of the automorphism induced by λ . The *Hilbert-Mumford weight* $\mu_H(\sigma, \lambda)$ determined by the polarization L , is the usual Hilbert-Mumford weight, that is, the weight of the $\mathbb{C}^\times$ -action on L over a generic value of $\text{Gr}(u)_0$.
- (c) (Ramanathan weight) Assume that $\lambda \in \mathfrak{r} \cong \mathfrak{r}^{\vee}$ is a weight of R . The Ramanathan weight of $\mu_R(\sigma, \lambda)$ of (P, u) with respect to (σ, λ) is given by the first Chern number of the line bundle determined by λ via the associated bundle construction: If $\mathbb{C}_\lambda$ is the one-dimensional representation of R with weight λ then the line bundle is $\sigma^*P \times_R \mathbb{C}_\lambda$ and the *Ramanathan weight* is

$$\mu_R(\sigma, \lambda) = \int_{[C]} c_1(\sigma^*P \times_R \mathbb{C}_\lambda).$$

- (d) (Mundet weight) The *Mundet weight* is the sum of the Hilbert-Mumford and Ramanathan weights:

$$\mu_M(\sigma, \lambda) := \mu_H(\sigma, \lambda) + \mu_R(\sigma, \lambda).$$

- (e) (Mundet semistability) The morphism (P, u) is *Mundet semistable* if

$$\mu_M(\sigma, \lambda) \leq 0$$

for all such pairs (σ, λ) [47], and *Mundet stable* if the above inequalities are satisfied strictly.

More generally, a *nodal n -marked gauged map* consists of an n -marked nodal projective curve $(\hat{C}, \underline{z})$ and a morphism $u : \hat{C} \rightarrow C \times X/G$ of class $(1, d) \in H_2(C, \mathbb{Z}) \times H_2^G(X, \mathbb{Z})$, consisting of a principal component $C_0 \subset \hat{C}$ which maps under u and projection to the first factor isomorphically to C , and a collection of *bubble components* $C_1, \dots, C_k$. The datum $(\hat{C}, u, \underline{z})$ is *Mundet semistable* if the restriction $u|_{C_0} : C_0 \rightarrow X/G$ is Mundet semistable, and each bubble component $C_i, i = 1, \dots, k$ on which u is given by a trivial G -bundle with constant section has at least three special (marked or nodal) points.

Let $\overline{\mathcal{M}}_n^G(C, X, d)$ denote the stack of Mundet semistable n -marked gauged maps. Taking X and G to be points, and d to be trivial, one obtains the *Fulton-MacPherson space* associated to C , that is, the moduli space

$$\overline{\mathcal{M}}_n(C) := \overline{\mathcal{M}}_{\text{genus}(C), n}(C, [C])$$

of stable maps to C of class $[C] \in H_2(C, \mathbb{Z})$. We denote by $\overline{\mathfrak{M}}_n(C)$ the stack of pre-stable maps of class $[C]$, that is, without the stability condition.

Proposition 1.4. *The category of Mundet-semistable gauged maps from C to X/G of homology class d and n markings forms an Artin stack $\overline{\mathcal{M}}_n^G(C, X, d)$ [25, 55], which if stable=semistable is a proper smooth Deligne-Mumford stack with perfect relative obstruction theory over $\overline{\mathfrak{M}}_n(C)$, evaluation map $\text{ev} : \overline{\mathcal{M}}_n^G(C, X, d) \rightarrow (X/G)^n$, and forgetful morphism $f : \overline{\mathcal{M}}_n^G(C, X, d) \rightarrow \overline{\mathcal{M}}_n(C)$.*

Example 1.5. (Toric Case) Suppose that G is a torus acting on a vector space X with weight decomposition $\bigoplus_{i=1}^k X_i$ so that G acts on the one-dimensional representation X_i with weights $\mu_i \in \mathfrak{g}^\vee$. Suppose that X is equipped with a polarization given by a trivial line bundle with character $\chi \in \mathfrak{g}^\vee$. Let C be a curve of genus 0. Then

$$\begin{aligned} \overline{\mathcal{M}}_0^G(C, X, d) &= \mathcal{M}_0^G(C, X, d) = H(\mathcal{O}_C(d) \times_G X) // G \\ &\cong \bigoplus_{i=1}^k X_i^{\oplus \langle \mu_i d \rangle + 1} // G \end{aligned}$$

where the polarization on $H^0(\mathcal{O}_C(d) \times_G X)$ is the trivial line bundle with character χ , see [55]. For example, if $G = \mathbb{C}^\times$ acts on $X = \mathbb{C}^k$ by scalar multiplication then $\overline{\mathcal{M}}_0^G(C, \mathbb{C}^k, d) = \mathbb{P}^{(d+1)k-1}$ for $d \geq 0$ and positive character χ .

Definition 1.6. (Gauged Gromov-Witten potential) Suppose that stable=semistable for all gauged maps. The gauged potential τ_X^G is the formal map defined by

$$\tau_X^G : QH_G(X) \rightarrow \Lambda_X^G, \quad \alpha \mapsto \sum_{n \geq 0, d \in H_2^G(X, \mathbb{Z})} (q^d/n!) \int_{[\overline{\mathcal{M}}_n^G(C, X, d)]} \text{ev}^*(\alpha, \dots, \alpha)$$

for $\alpha \in H_G(X)$, and extended to $QH_G(X)$ by linearity.

Remark 1.7. (Graph potential and gauged potential as quantizations of integration) Both $\tau_{X/G}$ and τ_X^G are quantizations of integration over the quotient in the following sense: Given $\alpha \in H_G(X)^n$ and a Fulton-MacPherson class $\beta \in H(\overline{\mathcal{M}}_n(C))$, consider the integral

$$(3) \quad \tau_{X/G}^n(\alpha, \beta) := \sum_d q^d \int_{[\overline{\mathcal{M}}_n(C, X/G, d)]} \text{ev}_n^* \alpha \cup f^* \beta.$$

If $\beta \in H^2(\overline{\mathcal{M}}_1(C)) \cong H^2(C)$ is the point class then

$$\tau_{X/G}^1(\alpha, \beta)|_{q=0} = \int_{[X/G]} \alpha.$$

Indeed any map from the curve C to $X//G$ of homology class 0 is constant. It follows that

$$\overline{\mathcal{M}}_n(C, X//G, 0) \cong (X//G) \times \overline{\mathcal{M}}_n(C).$$

In particular, for $n = 1$ we obtain $\overline{\mathcal{M}}_1(C, X//G, 0) \cong (X//G) \times C$, which implies the claim. A similar discussion holds for τ_X^G , that is,

$$\tau_{X/G}^1(\alpha, \beta)|_{q=0} = \int_{[X/G]} (D_0 \kappa_X^G|_{q=0}) \alpha.$$

The reader who is confused by the appearance of the Fulton-MacPherson class should keep in mind that a similar phenomenon appears in the Fukaya category setting: (co)units are defined by integrals over the moduli space of holomorphic disks with a single marking where the *position of the marking is fixed*. This ends the remark.

The gauged potential and the graph potential of the quotient are related by the *adiabatic limit theorem* of [55] (which is a generalization of an earlier result of Gaiotto-Salamon [23]): Let ρ be a real *stability parameter*; we consider Mundet stability with respect to the polarization L^ρ with $\rho \rightarrow \infty$.

Theorem 1.8 (Adiabatic limit theorem). *If $X//G$ is a locally free quotient then stable=semistable for ρ sufficiently large (more precisely, for any class $d \in H_2^G(X, \mathbb{Z})$ there exists an $r > 0$ such that $\rho > r$ implies stable=semistable) and*

$$\tau_{X/G} \circ \kappa_X^G = \lim_{\rho \rightarrow \infty} \tau_X^G.$$

In other words, the diagram

$$(4) \quad \begin{array}{ccc} QH_G(X) & \xrightarrow{\kappa_X^G} & QH(X//G) \\ & \searrow \tau_X^G \quad \swarrow \tau_{X/G} & \\ & \Lambda_X^G & \end{array}$$

commutes in the limit $\rho \rightarrow \infty$.

There is more general version of the adiabatic limit theorem for *Euler class twistings*: The universal curve $p : \overline{\mathcal{C}}_n^G(C, X) \rightarrow \overline{\mathcal{M}}_n^G(C, X)$ admits a universal gauged map $e : \overline{\mathcal{C}}_n^G(C, X) \rightarrow X/G$. For any G -equivariant bundle $E \rightarrow X$ we denote by

$$(5) \quad \text{Ind}(E) := Rp_*e^*(E/G)$$

the index of the bundle $E/G \rightarrow X/G$; this is an object in the bounded derived category of $\overline{\mathcal{M}}_n^G(C, X)$, since p is proper. The Euler class

$$(6) \quad \epsilon(E) := \text{Eul}_{\mathbb{C}^\times}(\text{Ind}(E)) \in H_{\mathbb{C}^\times}(\overline{\mathcal{M}}_n^G(C, X))$$

is well-defined after passing to the equivariant cohomology of $\overline{\mathcal{M}}_n^G(C, X)$ for the trivial $\mathbb{C}^\times$ -action corresponding to scalar multiplication on the fibers and inverting the equivariant parameter.

In order to study the dependence of the graph Gromov-Witten potential of the quotient on the choice of polarization, suppose that $L_\pm \rightarrow X$ are two polarizations of X and

$$L_t := L_-^{(1-t)/2} L_+^{(1+t)/2}$$

is the family of rational polarizations given by interpolation. Let $X//_\pm G$ denote the git quotients, $\kappa_{X,\pm}^G : QH_G(X, L_\pm) \rightarrow QH(X//_\pm G)$ the *quantum* Kirwan maps (note that we do not introduce new notation for the quantum version; the classical Kirwan map is obtained by setting $q = 0$) and $\tau_{X//_\pm G} : QH(X//_\pm G) \rightarrow \Lambda_{X,L_\pm}^G$ the graph potentials. The main result of this note is a formula for the difference $\tau_{X//_+ G} \kappa_{X,+}^G - \tau_{X//_- G} \kappa_{X,-}^G$ as a sum of fixed point contributions given by gauged Gromov-Witten invariants with smaller structure group. Namely, for any $t \in (-1, 1)$ such that $X^{G,t}$ is non-empty, there exists a stack $\overline{\mathcal{M}}_n^G(C, X, L_t)^G$ of G -fixed n -marked gauged maps from C to X/G consisting of a principal component mapping to $X^{G,t}/G$ and bubbles mapping to X/G . We denote by $\text{Ind}(T(X/G))^+$ the moving part of the normal complex $T(X/G)$ defined in (5), with Euler class

$$(7) \quad \epsilon_+(T(X/G)) := \text{Eul}_G(\text{Ind}(T(X/G))^+) \in H(\overline{\mathcal{M}}_n^G(C, X, L_t)^G)[\xi, \xi^{-1}].$$

The Mundet stability condition in general depends on a choice of equivariant Kähler class; here we are interested in the *large area limit* in which the gauged Gromov-Witten invariants are related to the git quotient. We denote by

$$\tilde{\Lambda}_X^G := \text{Hom}(H_2^G(X, \mathbb{Z}), \mathbb{Q})$$

the space of $\mathbb{Q}$ -valued functions on $H_G^2(X, \mathbb{Z})$. Note that $\tilde{\Lambda}_X^G$ has no ring structure extending that on Λ_X^G . The space $\tilde{\Lambda}_X^G$ can be viewed as the space of distributions in the quantum parameter q .

Definition 1.9 (Fixed point potential). Let $X, G, L_\pm$ be as above, and $t \in (-1, 1)$ such that $X^{G,t}$ is non-empty. The *fixed point potential* associated to this data is the map

$$(8) \quad \tau_{X,G,t}^G : QH_{G,\text{fin}}(X) \rightarrow \tilde{\Lambda}_X^G[\xi, \xi^{-1}]$$

$$\alpha \mapsto \sum_{d \in H_2^G(X, \mathbb{Z})} \sum_{n \geq 0} \int_{[\overline{\mathcal{M}}_n^G(C, X, L_t, d)^G]} (q^d/n!) \text{ev}^*(\alpha, \dots, \alpha) \cup \epsilon_+(T(X/G))^{-1}$$

for $\alpha \in H_G(X)$, extended to $QH_{G,\text{fin}}(X)$ by linearity.

We may now state the main result of the paper, in the case of circle actions.

Theorem 1.10 (quantum Kalkman formula, circle group case). *Suppose that $G = \mathbb{C}^\times$ and X is a smooth projective G -variety equipped with polarizations $L_\pm \rightarrow X$, and stable=semistable for $L_\pm$. Then*

$$(9) \quad \tau_{X/-G} \kappa_{X,+}^G - \tau_{X/+G} \kappa_{X,-}^G = \sum_{t \in (-1,1)} \text{Resid}_\xi \tau_{X,G,t}^G.$$

In other words, failure of the following square to commute is measured by an explicit sum of wall-crossing terms given by certain gauged Gromov-Witten invariants:

$$\begin{array}{ccccc} QH_G(X, L_-) & \longleftarrow & QH_G^{\text{fin}}(X) & \longrightarrow & QH_G(X, L_+) \\ \downarrow \kappa_{X,-}^G & & & & \downarrow \kappa_{X,+}^G \\ QH(X//_-G) & & & & QH(X//_+G) \\ \downarrow \tau_{X/-G} & & & & \downarrow \tau_{X/+G} \\ \Lambda_{X,L_-}^G & \longrightarrow & \tilde{\Lambda}_X^G & \longleftarrow & \Lambda_{X,L_+}^G. \end{array}$$

The diagram is somewhat more complicated than in the classical case in (4), because the maps $\tau_{X/\pm G} \kappa_{X,\pm}^G$ are defined using different the Novikov rings $\Lambda_{X,L_\pm}^{G, \geq 0}$. If a symbol q^d appears in one Novikov ring $\Lambda_{X,L_\pm}^{G, \geq 0}$ but not the other then the corresponding contribution to $\tau_{X/\pm G} \kappa_{X,\pm}^G$ must be equal to the q^d -term in the wall-crossing contribution on the right. See Example 1.11.

The wall-crossing formula for Gromov-Witten invariants Theorem 1.10 should be considered mirror to various results in on the behavior of the derived category of bounded complexes of coherent sheaves under variation of git quotient appearing recently in Segal [51], Halpern-Leistner [29] and Ballard-Favero-Katzarkov [3], and

more generally for crepant birational transformations, earlier in [35]. More precisely, one would hope that these results have extensions to open-closed Gromov-Witten theory, in particular, semi-orthogonal decompositions of Fukaya categories similar to those for the derived categories of matrix factorizations in the works above. We have stated the formula (9) in its simplest form; there are various extensions which include:

- (a) (Insertions from Fulton-MacPherson spaces) The moduli stacks $\overline{\mathcal{M}}_n^G(C, X)$ resp. $\overline{\mathcal{M}}_n(C, X//G)$ admit *forgetful morphisms* f to the moduli stack $\overline{\mathcal{M}}_n(C)$ of homology class $[C]$ stable maps to C . For any class $\beta \in H(\overline{\mathcal{M}}_n(C))$, one may insert $f^*\beta$ in the integral defining the potentials as in (3). The wall-crossing formula (9) then holds in the sense of *homotopy compositions of traces and morphisms of CohFT algebras* discussed in [55]. In fact, without the Fulton-MacPherson insertions the potentials $\tau_{X/G}, \tau_X^G$ will often vanish, unless one interprets them equivariantly in the case of a curve C of genus zero with a circle action, in which case they are closely related to Givental's I and J functions [24].
- (b) (Fixed point potential as a quantization of Kalkman's) The fixed point potential $\tau_{X,G,t}^G$ quantizes the fixed point contributions in Kalkman's formula (2), in the following sense: Let C be a genus zero curve and consider the n -th Taylor coefficient derivative $\tau_{X,G,t}^{G,n}$. Consider the integral with insertion of the Fulton-MacPherson class $\beta \in H(\overline{\mathcal{M}}_n(C))$ given by

$$\tau_{X,G,t}^{G,n}(\alpha, \beta) := \sum_d q^d \int_{[\overline{\mathcal{M}}_n^G(C, X, L_t, d)^G]} \text{ev}_n^* \alpha \cup f^* \beta \cup \epsilon_+(T(X/G))^{-1}.$$

If $\beta \in H(\overline{\mathcal{M}}_1(C)) \cong H(C)$ is the point class then

$$\tau_{X,G,t}^{G,1}(\alpha, \beta)|_{q=0} = \int_{[X^{G,t}]} \alpha \cup \text{Eul}_G(\nu_{X^{G,t}})^{-1}$$

which appears in Kalkman's wall-crossing formula. Indeed any map from the genus zero curve C to X/G of homology class 0 consists of a trivial bundle and constant section. It follows that

$$\overline{\mathcal{M}}_n^G(C, X, L_t, 0)^G \cong X^{G,t} \times \overline{\mathcal{M}}_n(C).$$

In particular, for $n = 1$ we obtain $\overline{\mathcal{M}}_1^G(C, X, L_t, 0)^G \cong X^{G,t} \times C$, which implies the claim. Furthermore, the Euler class $\epsilon_+(T(X/G))$ is the Euler class of the normal bundle to $X^{G,t}$.

- (c) (Twistings by Euler classes) One can introduce *twisted Gromov-Witten invariants* as follows. For any equivariant bundle E on X , inserting the Euler class $\epsilon(E) = \text{Eul}(\text{Ind}(E/G))$ of (6) gives twisted gauged Gromov-Witten invariants. Introducing similar twistings in the definition of κ_X^G , the wall-crossing formula extends to this case as well.

- (d) (Wall-crossing for individual Gromov-Witten invariants) Although we have written the formula as a difference of potentials, after unraveling the definitions one obtains wall-crossing formulas for individual Gromov-Witten invariants, or at least finite combinations of them. In Fano cases, one can often obtain wall-crossing formulas for individual Gromov-Witten invariants.
- (e) (Wall-crossing for non-convex actions) In some cases, even though the action of G on X is not convex at infinity (e.g. G is a torus acting on a vector space X with weights not contained in any half-space) the moduli spaces $\overline{\mathcal{M}}_n^G(C, X)$ are non-compact, but the moduli spaces admit an auxiliary group action with proper fixed point loci, and the maps $\tau_{X/G}, \kappa_X^G$ etc. can be defined via localization. The wall-crossing formula holds in this case as well, as long as the auxiliary group action extends to the various master spaces involved and the fixed point loci on these master spaces is proper (after fixing the homology class of the gauged map.) See Example 1.14 below.

Example 1.11. (Three-point Gromov-Witten invariants of projective space) The following simple example illustrates the notation involved in Theorem 1.10. As in Example 1.2, let $G = \mathbb{C}^\times$ acting on $X = \mathbb{C}^k$ by scalar multiplication, so that $H_G(X) = \mathbb{Q}[\xi]$. Suppose that polarizations $L_\pm$ correspond to the characters ± 1 , quotients $X//_-G = \emptyset$ and $X//_+G = \mathbb{P}^{k-1}$, and a unique singular value $t = 0$ corresponding to the point $0 \in X$. Let $\omega \in H^2(X//_+G)$ denote the hyperplane class. For integers $a, b, c \geq 0$ we compute genus 0 $d = 1$ $n = 3$ invariants $\langle \omega^a, \omega^b, \omega^c \rangle_{0,1}$ of $\mathbb{P}^{k-1}$ via wall-crossing; this was already covered in Cieliebak-Salamon [12]. Since $c_1^G(X) = k\xi$, the minimal Chern number of X is k . For dimensional reasons, there are no quantum corrections in $D_0\kappa_X^G(\xi^i), i \leq k-1$ and so $D_0\kappa_X^G(\xi^i) = \omega^i, i \leq k-1$. The adiabatic limit Theorem 1.8 with Fulton-MacPherson insertions implies that the genus zero, three-point invariants $\langle \omega^a, \omega^b, \omega^c \rangle_{0,d}$ of class $d \in H_2(X//G) \cong \mathbb{Z}$ in the quotient $X//G$ equal gauged Gromov-Witten invariants:

$$\langle \omega^a, \omega^b, \omega^c \rangle_{0,d} = \int_{[\overline{\mathcal{M}}_3^G(\mathbb{P}, X, d)]} \text{ev}_1^* \xi^a \cup \text{ev}_2^* \xi^b \cup \text{ev}_3^* \xi^c \cup f^* \beta$$

where $\mathbb{P}$ is the projective line and $\beta \in H^6(\overline{\mathcal{M}}_3(\mathbb{P}))$ is the point class, that is, the class fixing the location of the three marked points. Consider the wall-crossing formula Theorem 1.10, applied to the third order derivatives of the potential in the directions ξ^a, ξ^b, ξ^c with the Fulton-MacPherson insertion described above. There are no holomorphic spheres in X , so the moduli stack $\overline{\mathcal{M}}_0^G(\mathbb{P}, X, L_t, d)^G$ is a point, consisting of the bundle P with first Chern class $c_1(P) = d \in H_G^2(X, \mathbb{Z}) \cong \mathbb{Z}$ with constant section equal to zero. Thus the fixed point stack is

$$\overline{\mathcal{M}}_3^G(\mathbb{P}, X, L_t, d)^G \cong \overline{\mathcal{M}}_3(\mathbb{P}).$$

The index bundle of TX at the fixed point is

$$\text{Ind}(T(X/G))|_0 = H^0(\mathbb{P}, \mathcal{O}_{\mathbb{P}}(1) \times_{\mathbb{C}^\times} \mathbb{C}^k) \cong \mathbb{C}^{2k}.$$

It has Euler class $\epsilon_+(T(X/G)) = \xi^{2k}$. By the wall-crossing formula 1.10

$$\begin{aligned} \langle \omega^a, \omega^b, \omega^c \rangle_{0,1} &= \text{Resid}_\xi \int_{[\overline{\mathcal{M}}_3^G(\mathbb{P}, X, L_{0,1})^G]} \frac{\text{ev}_1^* \xi^a \cup \text{ev}_2^* \xi^b \cup \text{ev}_3^* \xi^c \cup f^* \beta}{\epsilon_+(T(X/G))} \\ &= \text{Resid}_\xi \xi^{a+b+c} / \xi^{2k} \\ &= \begin{cases} 1 & a+b+c = 2k-1 \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Thus

$$\omega^a \star \omega^b = q \omega^{a+b-k}, \quad k \leq a+b \leq 2k-1$$

as is well-known. For example, taking $a = b = k-1$ we obtain that there is a unique line in projective space passing through two generic points and a generic hyperplane. We give another Fano example in Example 3.7, where we compute the change in a coefficient in the fifth quantum power of the first Chern class for a blow-up of the projective plane. This ends the example.

The most interesting case of the wall-crossing formula Theorem 1.10 is the *crepant case* by which we mean that the sum of the weights at any fixed point vanishes (Definition 4.1); the term *crepant* was introduced by Reid [49] as the opposite of discrepant in the context of the minimal model program. In the last section we give a proof of a version of the crepant transformation conjecture of Li-Ruan [43], Bryan-Graber [10], Coates-Ruan [13] on equivalence of Gromov-Witten theories in this case:

Theorem 1.12 (Wall-crossing for crepant birational transformations of git type). *Suppose that X, G are as in Theorem 1.10, and C has genus zero. If all the wall-crossings are crepant then*

$$\tau_{X/-G} \circ \kappa_{X,-}^G \stackrel{a.e.}{=} \tau_{X/+G} \circ \kappa_{X,+}^G$$

almost everywhere (a.e.) in the quantum parameter q .

Remark 1.13. (a) Theorem 1.12 implies many of the special cases already known in the literature, although actually computing the transformations $\kappa_{X,\pm}^G$ relating the graph potentials $\tau_{X/\pm G}$ can be a difficult task. Note that Iwao-Lee-Lin-Wang [32], Lee-Lin-Wang [40] extend the invariance to cases not necessarily related by variation of git, while the results here are more general than that of [40], [32] since we allow “weighted flops”.

(b) Almost everywhere equality in the formal parameter q means the following: considering both sides as elements in $\text{Hom}(H_2^G(X, \mathbb{Z})/\text{torsion}, \mathbb{Q})$, the space of distributions on $H_G^2(X, \mathbb{R})/H_G^2(X, \mathbb{Z})$, the difference is a tempered distribution in at least one direction and its Fourier transform in that direction has support of measure zero, see Section 4.

(c) The proof of Theorem 1.12 uses an action of the Picard stack of the curve on the fixed point stacks. In the crepant case the (almost) invariance under this

action implies that, after summing over degrees, the wall-crossing term is a sum of derivatives of delta-functions in the quantum parameter q .

Example 1.14. (Simple three-fold flop, cf. Li-Ruan [43], Iwao-Lee-Lin-Wang [40], Lee-Lin-Wang [32]) The following simple example may help to explain the notion of “almost everywhere vanishing” of the wall-crossing contributions in the quantum parameter. Let $G = \mathbb{C}^\times$ acting on $X = \mathbb{C}^4$ with weights ± 1 each of multiplicity 2. Then

$$X//_{\pm}G = \mathcal{O}_{\mathbb{P}}(-1)^{\oplus 2},$$

where $\mathbb{P}$ is the projective line and we abuse notation by denoting by $\mathcal{O}_{\mathbb{P}}(-1)^{\oplus 2}$ the total space of two copies of the tautological line bundle $\mathcal{O}_{\mathbb{P}}(-1)$. Then $X//_+G$ and $X//_-G$ are isomorphic but the birational transformation relating them, induced by the variation of git quotient, is a simple flop. We consider the wall-crossing formula corresponding to the three-point invariants with each insertion given by $\xi \in H_G^2(X)$; that is, the third derivative of $\tau_{X//_{\pm}G}$ with a Fulton-MacPherson class $f^*\beta$ fixing the positions of the three marked points. In order to make sense of the non-compact integration in the $d = 0$ case one must take an equivariant extension of with respect to the action of $\mathbb{C}^\times$ acting by scalar multiplication, and use localization for the residual $\mathbb{C}^\times$ -action. Thus if θ is the equivariant parameter for the auxiliary $\mathbb{C}^\times$ then $\kappa_X^G(\xi + \theta) = \pm\omega + \theta$ where $\omega \in H^2(X//_{\pm}G)$ is the symplectic class integrating to 1 on the zero-section $\mathbb{P} \subset \mathcal{O}_{\mathbb{P}}(-1)^{\oplus 2}$. The fixed point set of the residual $\mathbb{C}^\times$ action is the zero section of $X//_{\pm}G$, which acts with weights ± 2 (with multiplicity two) on the normal bundle. The degree zero moduli space is $X//_{\pm}G$ itself, and integration over this non-compact space can be defined via localization as

$$\begin{aligned} I_{\pm} &:= \int_{X//_{\pm}G} \kappa_{X,\pm}^G(\alpha) := \int_{\mathbb{P}} (\pm\omega + \theta)^3 \cup \text{Eul}(\mathcal{O}_{\mathbb{P}}(-1)^{\oplus 2})^{-1} \\ &= \int_{\mathbb{P}} \frac{(\pm\omega + \theta)^3}{(-\omega \mp 2\theta)^{-2}} \\ &= \int_{\mathbb{P}} \pm 3\omega\theta^2(2\theta)^{-2} \pm \theta^3(2\theta)^{-2}(1 - \omega/(\pm 2\theta) + \dots)^2 \\ &= \pm 3/4 \mp 1/4 = \pm 1/2. \end{aligned}$$

The auxiliary circle action extends to the various master spaces involved, and the wall-crossing formula holds in this case as well, defining each contribution via localization. There is a unique fixed point $0 \in X$, with normal bundle isomorphic to X . Thus the wall-crossing term is

$$\text{Resid}_{\xi} \xi^3/\xi^2(-\xi)^2 = 1.$$

Thus Kalkman’s wall-crossing formula reduces to

$$I_+ - I_- = (1/2) - (-1/2) = 1.$$

Moving on to non-zero classes $d \in H_2^G(X, \mathbb{Z}) \cong \mathbb{Z}$, there is a unique gauged map of class d mapping to the fixed point $0 \in X$, with normal complex having weight 1

with multiplicity $2 + 2d$ and weight -1 with multiplicity $2 - 2d$, by Riemann-Roch. Thus the wall-crossing term class d is

$$\text{Resid}_\xi \xi^3 / \xi^{2+2d} (-\xi)^{2-2d} = 1.$$

The class d occurs in the expression $\tau_{X/\pm G} \kappa_{X,\pm}^G$ iff $d \geq 0$ resp. $d \leq 0$. Indeed, the contribution of q^d is the contribution of q^d to the gauged Gromov-Witten potential in Theorem 1.8, which by definition is an integral over Mundet stable maps which in this case is the git quotient of $H^0(\mathcal{O}_C(d) \times_{\mathbb{C}^\times} (\mathbb{C}_1^{\oplus d} \oplus \mathbb{C}_{-1}^{\oplus d}))$ by the $\mathbb{C}^\times$ action corresponding to the polarization $L_\pm$, where $\mathbb{C}_{\pm 1}$ denote the one-dimensional representations with weight ± 1 . For $d \neq 0$, only one factor $H^0(\mathcal{O}_C(d) \times_{\mathbb{C}^\times} (\mathbb{C}_{\pm 1}^{\oplus d})) \cong \mathbb{C}_{\pm 1}^{\oplus(d+1)}$ is non-vanishing and the git quotient is empty for the polarization $L_\mp$ corresponding to the character ∓ 1 . Since each $q^d, d \neq 0$ appears in only one of the Novikov rings $\Lambda_{X,\pm}^{G,\geq 0}$ we must have, after composing with the quantum Kirwan map, that the higher degree integrals for L_+ are 1 for $d > 0$ and 0 for $d < 0$ resp. for L_- are -1 for $d < 0$ and 0 for $d > 0$, each corresponding to an integral over multiple covers of the zero section. Summing over classes d the wall-crossing formula in Theorem 1.10 becomes

$$\left(\frac{1}{2} + \sum_{d>0} q^d \right) - \left(-\frac{1}{2} - \sum_{d<0} q^d \right) = \sum_{d \in \mathbb{Z}} q^d \stackrel{a.e.}{=} 0.$$

The reader may compare with the treatment of simple flops in [43], [40, Corollary 3.2] which contains essentially the same computation. Note that here we have not given an explicit description of the maps $\kappa_{X,\pm}^G$ which relate the two graph potentials. This ends the example.

2. KALKMAN'S WALL-CROSSING FORMULA

In this section we give a proof of Kalkman's formula 1.1 and give a generalization to non-abelian quotients.

2.1. The wall-crossing formula for circle actions. The wall-crossing formula is somewhat simpler for the case of a circle group, so we begin with that case. Let $G = \mathbb{C}^\times$ and X a smooth projective G -variety as above, equipped with polarizations $L_\pm \rightarrow X$. The proof of the wall-crossing formula is by localization on a proper Deligne-Mumford stack $\tilde{X}$ whose fixed points include $X//_\pm G$ and the fixed point components $X^{G,t}$ with $t \in (-1, 1)$. From the point of view of symplectic geometry, this is the *symplectic cut* construction in Lerman [41], but we need the algebraic approach here given in Thaddeus [52, Section 3].

First recall from the introduction that $L_t := L_-^{(1-t)/2} L_+^{(1+t)/2}$, $t \in (-1, 1)_\mathbb{Q}$ denotes the family of rational polarizations interpolating between $L_\pm$. We denote by $X//_t G$ the corresponding git quotients, by which we mean the stack-theoretic quotient of the semistable locus for L_t by the action of G . This means that if $\Phi_\pm$ are moment maps for action of a maximal compact of G on $L_\pm$ with respect to a unitary connection then $\Phi_t := (1-t)\Phi_-/2 + (1+t)\Phi_+/2$ is a moment map for the action of G on L_t . Even more concretely, if $\Phi_\pm$ are equal up to a constant $c = \Phi_+ - \Phi_-$ then

$\Phi_t = (\Phi_- + \Phi_+)/2 + tc/2$ are all equal up to a constant depending on t . For the following see Thaddeus [52, Section 3].

Lemma 2.1 (Existence of a master space). *Suppose that stable=semistable for $L_\pm$ and for any $t \in (-1, 1)$ and any t -semistable point $x \in X^G$, G_x acts with finite stabilizer on the fiber $(L_+ L_-^{-1})_x$. Then there exists a smooth proper Deligne-Mumford G -stack $\tilde{X}$ equipped with a line bundle ample for the coarse moduli space whose git quotients $\tilde{X} //_t G$ are isomorphic to those $X //_t G$ of X by the action of G with respect to the polarization L_t and whose fixed point set $\tilde{X}^G$ is given by the union*

$$\tilde{X}^G = X //_- G \cup X //_+ G \cup \bigcup_{t \in (-1, 1)} X^{G, t}.$$

where $X^{G, t}$ is the component of X^G that is semistable for parameter t .

Proof. The projectivization $\mathbb{P}(L_- \oplus L_+)$ of the direct sum $L_- \oplus L_+$ of the polarizations $L_\pm \rightarrow X$ has a natural polarization given by the relative hyperplane bundle $\mathcal{O}_{\mathbb{P}(L_- \oplus L_+)}(1)$ of the fibers, with stalks

$$(10) \quad \mathcal{O}_{\mathbb{P}(L_- \oplus L_+)}(1)_{[l_-, l_+]} = \text{span}(l_- + l_+)^{\vee}.$$

Let $\pi : \mathbb{P}(L_- \oplus L_+) \rightarrow X$ denote the projection. (A word of warning: the notation π will be used for a number of different projections in this paper.) The space of sections of $\mathcal{O}_{\mathbb{P}(L_- \oplus L_+)}(k)$ has a decomposition under the natural $\mathbb{C}^\times$ -action with eigenspaces given by the sections of $\pi^* L_-^{k_-} \oplus \pi^* L_+^{k_+}$ with $k_- + k_+ = k, k_\pm \geq 0$. It follows that the semistable locus in $\mathbb{P}(L_- \oplus L_+)$ is the union of $\pi^{-1}(X^{\text{ss}, t})$ where $X^{\text{ss}, t} \subset X$ is the semistable locus for $L_-^{(1-t)/2} L_+^{(1+t)/2}, t \in [-1, 1]$. Let $\tilde{X} := \mathbb{P}(L_- \oplus L_+) // G$ denote the geometric invariant theory quotient, by which we mean the stack-theoretic quotient of the semistable locus. The assumption on the action of the stabilizers implies that the action of G on the semistable locus in $\mathbb{P}(L_- \oplus L_+)$ is locally free, so that stable=semistable for $\mathbb{P}(L_- \oplus L_+)$. It follows that $\tilde{X}$ is a proper smooth Deligne-Mumford stack. The quotient $\tilde{X}$ contains the quotients of $\mathbb{P}(L_\pm) \cong X$ with respect to the polarizations $L_\pm$, that is, $X //_\pm G$. The G -action with weights ± 1 on $L_- \oplus L_+$ induces a G -action on $\tilde{X}$ with fixed point set as described above. Indeed, the fixed points are pairs $[l_-, l_+]$ such that either $l_- = 0, l_+ = 0$, or l_-, l_+ are both non-zero but the projection to X is G -fixed. $\square$

Remark 2.2. (Condition for stable=semistable) The proof shows that the G -action on $\mathbb{P}(L_- \oplus L_+)$ has stable=semistable iff the condition of the lemma holds, that is, stable=semistable for $L_\pm$ and for any $t \in (-1, 1)$ and t -semistable point $x \in X^G$, G_x acts with finite stabilizer on the fiber $(L_+ L_-^{-1})_x$. Indeed, this is the condition for G to act locally freely on the semistable locus in $\mathbb{P}(L_- \oplus L_+)$.

The projection $\mathbb{P}(L_- \oplus L_+) \rightarrow X$ induces a G -invariant map $\tilde{X} \rightarrow X$. Composing with the Kirwan map one obtains a canonical map $H_G(X) \rightarrow H_G(\tilde{X})$. Its composition with $\tilde{\kappa}_\pm : H_G(\tilde{X}) \rightarrow H(X //_\pm G)$ is equal to $\kappa_{X, \pm}^G$.

Proof of Kalkman's wall-crossing Theorem 1.1. Taking the residue of the localization formula for the G -action on $\tilde{X}$ gives Theorem 1.1, see Lerman [41]. Indeed for any equivariant class $\alpha \in H_G(X)$ of top degree, its pullback to $\mathbb{P}(L_- \oplus L_+)$ descends to a class $\tilde{\alpha} \in H_G(\tilde{X})$ whose restriction to $X//_{\pm}G$ is $\kappa_{X,\pm}^G(\alpha)$, and whose restriction to $X^{G,t} \subset \tilde{X}^G$ is $\iota_{X^{G,t}}^* \alpha$. Since $\deg(\tilde{\alpha}) = \deg(\alpha) = \dim(\tilde{X}) - 2$, the integral of $\tilde{\alpha}$ over $\tilde{X}$ vanishes. On the other hand, by localization the integral is

$$\int_{[\tilde{X}]} \alpha = \int_{[X//_G]} \frac{\kappa_{X,-}^G(\alpha)}{\text{Eul}_G(\nu_-)} + \int_{[X//_+G]} \frac{\kappa_{X,+}^G(\alpha)}{\text{Eul}_G(\nu_+)} + \sum_{t \in (-1,1)} \int_{[X^{G,t}]} \frac{\iota_{X^{G,t}}^* \alpha}{\text{Eul}_G(\nu_{X^{G,t}})}$$

where $\nu_{\pm} \rightarrow X//_{\pm}G$ are the normal bundles to $X//_{\pm}G$ in $\tilde{X}$. The inverted Euler classes are

$$\text{Eul}_G(\nu_{\pm})^{-1} = (\mp \xi + c_1(\nu_{\pm}))^{-1} = \mp \xi^{-1}(1 + c_1(\nu_{\pm})(\mp \xi)^{-1} + \dots).$$

Hence taking residues one obtains

$$0 = -\tau_{X//_+G} \kappa_{X,+}^G \alpha + \tau_{X//_G} \kappa_{X,-}^G \alpha + \text{Resid}_{\xi} \sum_{t \in (-1,1)} \int_{[X^{G,t}]} \frac{\iota_{X^{G,t}}^* \alpha}{\text{Eul}_G(\nu_{X^{G,t}})}$$

as claimed. $\square$

2.2. Kalkman wall-crossing for actions of non-abelian groups. Kalkman's wall-crossing formula can be used to study the intersection pairings for variation of git for the action of an arbitrary connected complex reductive group G on a smooth polarized projective variety X as follows. Let us examine the structure of the fixed point set of $\mathbb{C}^{\times}$ on $\tilde{X}$. For any $\zeta \in \mathfrak{g}$, we denote by $G_{\zeta} \subset G$ the stabilizer of ζ under the adjoint action of G .

Let $T \subset G$ be a maximal torus and $\mathfrak{t} \subset \mathfrak{g}$ the corresponding Cartan algebra. Let $W = N(T)/T$ denote the Weyl group. For any $\zeta \in \mathfrak{t}$, we denote by W_{ζ} resp. $W_{\mathbb{C}\zeta}$ the group of $w \in W$ that fix the element $\zeta \in \mathfrak{g}$ resp. line $\mathbb{C}\zeta \subset \mathfrak{g}$. Thus the quotient $W_{\mathbb{C}\zeta}/W_{\zeta}$ is either isomorphic to $\{\pm 1\}$ or to $\{1\}$, depending on whether or not there is a Weyl group element acting as -1 on $\mathbb{C}\zeta$.

Lemma 2.3 (Structure of the fixed point components). *Suppose that stable=semistable for the G -action on $\mathbb{P}(L_- \oplus L_+)$, so that $\tilde{X}$ is a smooth proper Deligne-Mumford stack with $\mathbb{C}^{\times}$ action, constructed in the proof of Lemma 2.1.*

- (a) Any fixed point $\tilde{x} \in \tilde{X}^{\mathbb{C}^{\times}}$ with $\tilde{x} = [l]$ for some $l \in \mathbb{P}(L_- \oplus L_+)$ has the property that for all $z \in \mathbb{C}^{\times}$, $z l = z^{\zeta} l$ for some unique $\zeta \in \mathfrak{g}$.
- (b) For each $\zeta \in \mathfrak{t}$ there is a morphism $\iota_{\zeta} : X^{\zeta} //_{\mathfrak{t}} (G_{\zeta} / \mathbb{C}^{\times}) \rightarrow \tilde{X}^{\mathbb{C}^{\times}}$ with fiber $W_{\mathbb{C}\zeta} / W_{\zeta}$. The images of ι_{ζ} cover $\tilde{X}^{\mathbb{C}^{\times}}$, disjointly after passing to equivalence classes of one-parameter subgroups.
- (c) For any $\alpha \in H_G(X)$, the pull-back of $\tilde{\kappa}(\alpha)|_{\tilde{X}^{\mathbb{C}^{\times}}}$ under ι_{ζ} is equal to image of α under the restriction map $H_G(X) \rightarrow H_{\mathbb{C}^{\times}}(X^{\zeta} //_{\mathfrak{t}} (G_{\zeta} / \mathbb{C}^{\times}))$.
- (d) The pull-back of the normal bundle $N_{\tilde{X}^{\mathbb{C}^{\times}}}$ of $\tilde{X}^{\mathbb{C}^{\times}}$ under ι_{ζ} is canonically isomorphic to the image of $N_{X^{\zeta}} / (\mathfrak{g} / \mathbb{R}\zeta)$ under the quotient map $X^{\zeta} \rightarrow X^{\zeta} // (G_{\zeta} / \mathbb{C}^{\times})$,

by an isomorphism that intertwines the action of $\mathbb{C}_\zeta^\times$ on $(N_{X^\zeta}/(\mathfrak{g}/\mathbb{R}\zeta))// (G_\zeta/\mathbb{C}_\zeta^\times)$ with the action of $\mathbb{C}^\times$ on $N_{\tilde{X}^{\mathbb{C}^\times}}$.

Proof. If $[l] \in \tilde{X}^{\mathbb{C}^\times}$, with $l \in \mathbb{P}(L_- \oplus L_+)$ then $[l] \in \tilde{X}^\xi$, where ξ is a generator of the Lie algebra of $\mathbb{C}^\times$ and $\tilde{X}^\xi$ is the zero set of the vector field $\xi_{\tilde{X}}$ generated by ξ . Since $\tilde{X}$ is the quotient of $\mathbb{P}(L_- \oplus L_+)$ by G , if ξ_L denotes the vector field on $\mathbb{P}(L_- \oplus L_+)$ generated by ξ then $\xi_L(l) = \zeta_L(l)$ for some $\zeta \in \mathfrak{g}$. Since G acts locally freely ζ must be unique. Integrating gives $zl = z^\zeta l$ for all $z \in \mathbb{C}^\times$. By uniqueness of ζ , this holds for every point in the component of $\tilde{X}^\xi$ containing $[l]$. The fiber of the morphism $X^\zeta// (G_\zeta/\mathbb{C}_\zeta^\times) \rightarrow \tilde{X}^{\mathbb{C}^\times}$ is the set of $g \in G/G_\zeta$ fixing both G_ζ and $\mathbb{C}_\zeta^\times$ under the adjoint action. The set of $g \in G/G_\zeta$ fixing G_ζ is $N(G_\zeta)/G_\zeta \cong W/W_\zeta$, while the set of $g \in G/G_\zeta$ fixing $\mathbb{C}_\zeta^\times$ and G_ζ is isomorphic to $W_{\mathbb{C}_\zeta}/W_\zeta$. Hence the fiber is $W_{\mathbb{C}_\zeta}/W_\zeta$. The remaining claims are left to the reader. $\square$

The group $\mathbb{C}_\zeta^\times \subset G_\zeta$ acts trivially on X^ζ , which is therefore a $G_\zeta/\mathbb{C}_\zeta^\times$ -space. This is the “smaller structure group” referred to in the abstract. For any fixed point component $X^{\zeta,t} \subset X^\zeta$ that is t -semistable, we denote by $\nu_{X^\zeta,t}$ the normal bundle of $X^{\zeta,t}$ modulo $\mathfrak{g}/\mathbb{C}_\zeta$, quotiented by $G_\zeta/\mathbb{C}_\zeta^\times$. Let

$$\tau_{X^\zeta,t}^{G_\zeta} : H_G(X) \rightarrow \mathbb{Q}[\xi, \xi^{-1}], \quad \alpha \mapsto \int_{[X^{\zeta,t}/(G_\zeta/\mathbb{C}_\zeta^\times)]} \alpha \cup \text{Eul}_{\mathbb{C}_\zeta^\times}(\nu_{X^\zeta,t})^{-1}$$

where we have omitted the map $H_G(X) \rightarrow H_{\mathbb{C}_\zeta^\times}(X^{\zeta,t})$ to simplify notation, and ξ is the equivariant parameter for $\mathbb{C}_\zeta^\times$.

Theorem 2.4 (Kalkman wall-crossing). *Let X be a smooth projective G -variety and $L_\pm \rightarrow X$ polarizations with stable=semistable for the G action on $\mathbb{P}(L_- \oplus L_+)$. Then*

$$(11) \quad \tau_{X/+G} \kappa_{X,+}^G - \tau_{X/-G} \kappa_{X,-}^G = \sum_{t \in (-1,1), [\zeta]} \frac{|W_\zeta|}{|W_{\mathbb{C}_\zeta}|} \text{Resid}_\xi \tau_{X^\zeta,t}^{G_\zeta}$$

where the sum over $[\zeta]$ represents a sum over one-parameter subgroups of G , up to conjugacy.

Example 2.5. (a) (Blow-up of the projective plane as a quotient of a product of projective lines by a circle action) Let $X = (\mathbb{P})^3$ with polarization $\mathcal{O}_\mathbb{P}(a) \boxtimes \mathcal{O}_\mathbb{P}(b) \boxtimes \mathcal{O}_\mathbb{P}(c)$ with $a \ll b \ll c$. Let $\alpha = c_1^G(X)^2$. Since TX is the same as $T(X//_\pm G)$ plus a trivial line bundle, we have $\kappa_{X,\pm}^G(c_1^G(X)) = c_1(X//_\pm G)$. Furthermore, since all weights are 1, we have $c_1^G(X) = 3\xi$. Consider the wall-crossing from the chamber $\Phi < -a - b - c$ to the first non-empty chamber $\Phi \in (a - b - c, -a + b - c)$. There is a unique fixed point with weights 1, 1, 1 on the normal bundle; hence

$$\tau_{X/+G} \kappa_{X,+}^G(\alpha) - \tau_{X/-G} \kappa_{X,-}^G(\alpha) = \text{Resid}_\xi(3\xi)^2/\xi^3 = 9.$$

Indeed, $c_1(\mathbb{P}^2)$ is three times the generator of $H^2(\mathbb{P}^2)$, so

$$\tau_{X,-}^G \kappa_{X,-}^G(\alpha) = \int_{[\mathbb{P}^2]} c_1(\mathbb{P}^2)^2 = 9.$$

Consider next the wall-crossing from the chamber with quotient $X//_-G = \mathbb{P}^2$ to the chamber with quotient $X//_+G = \text{Bl}(\mathbb{P}^2)$, where $\text{Bl}(\mathbb{P}^2)$ is the blow-up of $\mathbb{P}^2$ at a point. Letting $\pi : \text{Bl}(\mathbb{P}^2) \rightarrow \mathbb{P}^2$ denote the blow-down map we have $c_1(\text{Bl } \mathbb{P}^2) = 3\pi^*H + E$ where H, E are the hyperplane class and class of the exceptional divisor respectively. Hence

$$\begin{aligned} \tau_{X//_+G} \kappa_{X,+}^G(\alpha) &= (3\pi^*H + E)^2 \\ &= (3\pi^*H)^2 + 6\pi^*HE + E^2 = 9 + 0 - 1 = 8. \end{aligned}$$

To compare this result with the wall-crossing formula, note that the fixed point set $X^{G,t}$ consists of a unique point $([0, 1], [1, 0], [1, 0])$ which is semistable exactly for $t = a - b - c$, with tangent weights $-1, 1, 1$. It follows that the first Chern class squared and Euler classes are

$$c_1(TX|X^{G,t})^2 = (-\xi + \xi + \xi)^2 = \xi^2, \quad \text{Eul}_G(TX|X^{G,t}) = -\xi^3.$$

Hence the wall-crossing term is

$$\text{Resid}_\xi \left(\int_{[X^{G,t}]} \iota_{X^{G,t}}^* \alpha \cup \text{Eul}_G(\nu_{X^{G,t}})^{-1} \right) = \text{Resid}_\xi \frac{\xi^2}{-\xi^3} = -1.$$

The wall-crossing formula reduces to $8 - 9 = -1$. In fact each blow-up of $\mathbb{P}^2$ lowers c_1^2 by 1, leading to the 9 del Pezzo surfaces, each of which contains a simple simply-laced root system corresponding to classes of self-intersection -2 etc. [44].

- (b) (Blow-up of the projective plane as a quotient of affine four-space by a two-torus) Let us do the same example in a different way, namely as a quotient of an affine space. $X = \mathbb{C}^4$ with $G = (\mathbb{C}^\times)^2$ acting with weights $(1, 0), (1, 0), (1, 1), (0, 1)$. Consider the path of polarizations in $H_G^2(X, \mathbb{Q}) \cong \mathbb{Q}^2$ from $(1, 2)$ to $(2, 1)$ so that $X//_-G$ is isomorphic to $\mathbb{P}^2$ via the map

$$X//_-G \rightarrow \mathbb{P}^2, \quad [a, b, c, d] = [(a, b, cd^{-1}, 1)] \mapsto [a, b, cd^{-1}].$$


On the other hand $X//_+G$ is isomorphic to the blow-up of $\mathbb{P}^2$ with the map to $\mathbb{P}^2$ blowing down the exceptional divisor given by

$$X//_+G \rightarrow \mathbb{P}^2, \quad [a, b, c, d] \mapsto [a, b, cd^{-1}].$$

As one moves in a line from say $(-1, 2)$ to $(2, -1)$ the symplectic quotients are in order $\emptyset, \mathbb{P}^2, \text{Bl}(\mathbb{P}^2), \emptyset$ where the initial and final contractions are projective bundles over the fixed point sets pt resp. $\mathbb{P}$ for the residual action of $\mathbb{C}^\times$ on the quotient of X by the diagonal action. See Figure 1, where the two chambers for the possible quotients are shown together with the moment polytopes of the quotients in each chamber.

Consider first the wall-crossing from the empty chamber to the non-empty chamber with quotient $\mathbb{P}^2$ in Figure 1. We have $c_1^G(X) = 3\xi_1 + 2\xi_2$. Hence the wall-crossing term is

$$\text{Resid}_{\xi=\xi_1} \frac{(3\xi_1 + 2\xi_2)^2}{\xi_1^2(\xi_1 + \xi_2)} \Big|_{\xi_2=0} = 9$$

FIGURE 1. Chambers for $(\mathbb{C}^\times)^2$ action on $\mathbb{C}^4$

hence $\int_{[\mathbb{P}^2]} c_1(\mathbb{P}^2)^2 = 9$ as expected.

The wall-crossing term for passing from the chamber with quotient $\mathbb{P}^2$ to that with quotient $\text{Bl}(\mathbb{P}^2)$ is

$$\text{Resid}_\xi \left. \frac{(3\xi_1 + 2\xi_2)^2}{\xi_1^2 \xi_2} \right|_{\xi=\xi_1=-\xi_2} = -1.$$

This implies that $\int_{[\text{Bl}(\mathbb{P}^2)]} c_1(\mathbb{P}^2)^2 = 8$ as we already computed in item (a) above.

Passing from the chamber with quotient $\text{Bl}(\mathbb{P}^2)$ to the adjacent chamber with empty quotient, note that the symplectic quotient at the wall-crossing point $\mu = (1, 0)$ equal to $\mathbb{P}$ and a wall-crossing term of

$$\text{Resid}_\xi \int_{[\mathbb{P}]} \kappa_X^{\mathbb{C}^\times} \frac{(3\xi_1 + 2\xi_2)^2}{\xi_2(\xi_1 + \xi_2)}$$

where $\kappa_X^{\mathbb{C}^\times} : H_G(X) \rightarrow H_{G/\mathbb{C}^\times}(X//\mathbb{C}^\times)$ is the Kirwan map for the quotient by the first factor of $\mathbb{C}^\times$, so that $X//\mathbb{C}^\times$ has $\mathbb{C}^\times$ -quotients $X//_\pm G$. Under $\kappa_X^{\mathbb{C}^\times}$ we have that ξ_1 maps to the generator ω of $H^2(\mathbb{P})$ while ξ_2 maps to the parameter ξ for the residual $\mathbb{C}^\times$ -action. The wall-crossing term is

$$\begin{aligned} \text{Resid}_\xi \int_{[\mathbb{P}]} \frac{(3\omega + 2\xi)^2}{\xi(\omega + \xi)} &= \text{Resid}_\xi \int_{[\mathbb{P}]} \frac{(3\omega + 2\xi)^2}{\xi^2} (1 - \omega/\xi + \omega^2/\xi^2 - \dots) \\ &= \text{Resid}_\xi \int_{[\mathbb{P}]} 12\omega/\xi - 4\omega/\xi = -8. \end{aligned}$$

Thus as expected the wall-crossing formula reduces to $\int_\emptyset c_1(\emptyset) = 8 - 8 = 0$.

- (c) (Resolution of a crepant singularity) The following example illustrates the application of the wall-crossing to orbifolds, in which the integrals become rational, and also the non-compact case, in which the integrals must be defined via localization. Suppose that $X = \mathbb{C}^{k+1}$ and $G = \mathbb{C}^\times$ acts with weights 1 with multiplicity k and $-k$ with multiplicity 1, so that $X//_+ G = \mathbb{C}^k/\mathbb{Z}_k$ (by our

conventions, a stack with trivial canonical bundle) while $X//_G$ is isomorphic to the total space of $\mathcal{O}_{\mathbb{P}^{k-1}}(k) \rightarrow \mathbb{P}^{k-1}$. Let $\alpha = c_k^G(\mathbb{C}^{k+1})$ (the next-to-highest Chern class of $\mathbb{C}^{k+1}$) so that $\kappa_{X,\pm}^G(\alpha)$ is the top Chern class of $X//_{\pm}G$. Then

$$\tau_{X/+G}(\kappa_{X,+}^G(\alpha)) = \int_{[\mathbb{C}^k/\mathbb{Z}_k]} \text{Eul}(\mathbb{C}^k/\mathbb{Z}_k) = 1/k.$$

Indeed, interpreting this integral via localization using the $\mathbb{C}^\times$ -action given by scalar multiplication, there is a unique fixed point with stabilizer of order k which contributes $\text{Eul}(\mathbb{C}^k)/\text{Eul}(\mathbb{C}^k)k = 1/k$ to the integral. On the other side of the wall,

$$\tau_{X/-G}(\kappa_{X,-}^G(\alpha)) = \int_{[\mathcal{O}_{\mathbb{P}^{k-1}}(k) \rightarrow \mathbb{P}^{k-1}]} \text{Eul}(\mathbb{P}^{k-1}) = k.$$

The wall-crossing term is

$$\text{Resid}_\xi \frac{(1+\xi)^k(1-k\xi)}{\xi^k(-k\xi)} = k - 1/k$$

as expected.

- (d) (A del Pezzo surface as a quotient of a product of projective lines by a non-abelian group) Let $X = (\mathbb{P})^5$ with the diagonal action of $G = PGL(2, \mathbb{C}) \cong SO(3, \mathbb{C})$, equipped with the polarization $\mathcal{O}(1)^t \boxtimes \mathcal{O}(1)^{\boxtimes 4}$ corresponding to varying the symplectic class on the first factor. From the symplectic point of view, this is the symplectic quotient of $(S^2)^5$ by the $SO(3)$ -action with radii of the spheres $(t, 1, 1, 1, 1)$. The moment map for the action is

$$X \rightarrow \mathfrak{g}_{\mathbb{R}}^\vee = \mathfrak{so}(3)^\vee \cong \mathbb{R}^3, \quad (x_1, \dots, x_5) \mapsto x_1 + \dots + x_5.$$

The quotient $X//G$ is homeomorphic to the *moduli space of polygons*

$$X//G \cong \left\{ (x_1, \dots, x_5) \left| \sum_{j=1}^5 x_j = 0, \|x_1\| = t, \|x_j\| = 1, j > 1 \right. \right\} / SO(3).$$

A point in the zero level set of the moment map is fixed by a one-parameter subgroup iff the points $(x_1, \dots, x_5)$ are colinear, in which case the subgroup fixing them is rotation around the corresponding line. One sees easily from this (or the algebraic description) that the singular values of t are 2, 4 corresponding to a single orbit of colinear configurations $x_1 = x_2 + x_3 + x_4 = x_5$ when $\|x_1\| = t = 4$ and four orbits of colinear configurations $x_1 + x_i = x_j + x_k + x_l$ when $\|x_1\| = t = 2$. Thus the chamber structure for t is $(0, 2), (2, 4), (4, \infty)$. Since the quotient in the last chamber is empty, and all the weights are one, the quotient $X//_tG$ for the last chamber must be $\mathbb{P}^2$, while passing from the second to first chamber the quotient undergoes a blow-up at 4 points. Consider the wall-crossing given by passing between chambers so that $X//_G = \mathbb{P}^2$ and $X//_+G = \text{Bl}^4(\mathbb{P}^2)$. We compute the square of the first Chern class by wall-crossing: Let $\alpha = c_1^G(X)^2$, so that $\kappa_{X,\pm}^G(\alpha) = c_1(X//_{\pm}G)^2$. The fixed points of $\mathbb{C}^\times$ action for the given singular value are the 8 configurations with

$x_1 + x_i = x_j + x_k + x_l$ with all colinear. Consider the case that $x = (x_1, \dots, x_5)$ is fixed by the maximal torus action; the tangent bundle of this point has weights $1, 1, 1, -1, -1$ hence $c_1^G(X) = \xi$ at x , while the normal bundle modulo $\mathfrak{g}/\mathbb{C}\zeta$ has weights $1, 1, -1$. Taking into account the factor $|W_\zeta|/|W_{\mathbb{C}\zeta}| = 1/2$ (corresponding to the fact that the configurations related by multiplication by -1 have the same image in $\tilde{X}^G$) we have

$$\begin{aligned} \int_{[X/+G]} c_1(X//_+G)^2 &= \int_{[X/-G]} c_1(X//_-G)^2 + 4 \operatorname{Resid}_\xi \frac{\xi^2}{-\xi^3} \\ &= 9 - 4 = 5 \end{aligned}$$

which matches the fact that each blow-up of $\mathbb{P}^2$ lowers c_1^2 by 1.

2.3. The virtual wall-crossing formula. Our main result will be derived from a virtual extension of wall-crossing formulas, carried out by Kiem-Li [37] using the virtual localization formula of Graber-Pandharipande [27]; we assume that the reader is familiar with the concepts of equivariant perfect obstruction theories etc. from those papers. In this section we explain the virtual extension and give an application to a simple complete intersection.

Let $\mathcal{X}$ be a Deligne-Mumford G -stack equipped with a G -equivariant perfect obstruction theory which admits an embedding in a smooth Deligne-Mumford G -stack. This means that $\mathcal{X}$ is equipped with an object E of the bounded derived category of G -equivariant coherent sheaves, together with a morphism from E to the cotangent complex, satisfying certain axioms, see [4], [27]; a typical example is an invariant complete intersection as in Example 2.7 below. If $\zeta \in \mathfrak{g}$ is any element then the fixed point stacks $\mathcal{X}^\zeta$ also have equivariant perfect obstruction theories compatible with that on $\mathcal{X}$; we denote by $\nu_{\mathcal{X}^\zeta}$ the virtual normal complex for the embedding $\mathcal{X}^\zeta \rightarrow \mathcal{X}$. The Euler class $\operatorname{Eul}_{\mathbb{C}^\times}(\nu_{\mathcal{X}^\zeta})$ is well-defined in the equivariant cohomology $H_{\mathbb{C}^\times}(\mathcal{X}^\zeta)$, after inverting the equivariant parameter ξ . From the virtual localization formula one obtains the following virtual version of the wall-crossing Theorem 2.4, which can be considered a corollary of Kiem-Li [37]:

Theorem 2.6 (Virtual Kalkman wall-crossing). *Let $\mathcal{X}$ be a proper Deligne-Mumford G -stack equipped with a G -equivariant perfect obstruction theory which admits an embedding in a smooth Deligne-Mumford G -stack. Let $L_\pm \rightarrow \mathcal{X}$ be a G -line bundles that are ample for the coarse moduli spaces, so that the stable=semistable for $\mathbb{P}(L_- \oplus L_+)$. Let $\tau_{\mathcal{X}/\pm G}$ resp. $\tau_{\mathcal{X},\zeta,t}^{G_\zeta}$ denote integration resp. equivariant integration over $\mathcal{X}/\pm G$ resp. $\mathcal{X}^{\zeta,t}$ times $\operatorname{Eul}_{\mathbb{C}^\times}(\nu_{\mathcal{X}^{\zeta,t}})^{-1}$. Then*

$$(12) \quad \tau_{\mathcal{X}/+G} \kappa_{\mathcal{X},+}^G - \tau_{\mathcal{X}/-G} \kappa_{\mathcal{X},-}^G = \sum_{t \in (-1,1), [\zeta]} \frac{|W_\zeta|}{|W_{\mathbb{C}\zeta}|} \operatorname{Resid}_\xi \tau_{\mathcal{X},\zeta,t}^{G_\zeta}$$

where the sum over $[\zeta]$ represents a sum over unparametrized one-parameter subgroups of G , up to conjugacy.

Example 2.7. (Wall-crossing over a nodal fixed point) The following simple example may help illustrate the notation. Suppose that X is a nodal projective line with a

single node, equipped with the standard $\mathbb{C}^\times$ -action on each component, so that the weights of the action on the tangent spaces the node are ± 1 . We equip X with a polarization so that the weights are ± 1 at the smooth fixed points, and 0 at the nodal point. Then $X//_t G$ is a point for $t \in (-1, 1)$, and is singular for $t \in \{-1, 0, 1\}$. Since X is complete intersection, X is virtually smooth and the virtual wall-crossing formula of Theorem applies. We examine the wall-crossing for the trivial class $\alpha = 1$ at the singular value $t = 0$. The virtual normal complex at the nodal point is the quotient of $\mathbb{C}_1 \oplus \mathbb{C}_{-1}$, the sum of one-dimensional representations with weights 1, -1 , modulo their tensor product $\mathbb{C}_1 \otimes \mathbb{C}_{-1}$, which has weight zero. Hence the normal complex has inverted Euler class

$$\mathrm{Eul}_G(\nu_{X,G,t})^{-1} = \frac{0\xi}{\xi(-\xi)} = 0.$$

The integrals on the left and right hand sides are 1 (being the integrals over points) while the wall-crossing term is

$$\begin{aligned} 1 - 1 &= \tau_{X/+G} \kappa_{\mathcal{X},+}^G - \tau_{X/-G} \kappa_{\mathcal{X},-}^G \\ &= \mathrm{Resid}_\xi \tau_{\mathcal{X},\zeta,0}^G = \mathrm{Resid}_\xi \int_{[\mathrm{pt}]} \mathrm{Eul}_G(\nu_{X,G,t})^{-1} = \mathrm{Resid}_\xi 0 = 0 \end{aligned}$$

as desired.

3. WALL-CROSSING FOR GROMOV-WITTEN INVARIANTS

In this section, we prove a quantum generalization of Kalkman's wall-crossing formula Theorem 2.4. In the first two subsections, we define the wall-crossing terms as integrals over moduli spaces of gauged maps fixed by a central subgroup. The last two subsections contain a construction of a master space for moduli spaces of gauged maps, and a proof of the wall-crossing formula via virtual localization on the master space. The construction of the master space is obtained from one for a different compactification of gauged maps introduced by Schmitt [50], pulled back under a relative version of Givental's morphism from stable maps to the quot scheme. Schmitt's compactification has the advantage that it is constructed by git methods so the classical techniques apply.

3.1. Gauged maps for groups with center. Let G be a complex connected reductive group and $Z \subset G$ a central subgroup. For any principal G -bundle $P \rightarrow C$, the right action of Z induces an action on the associated bundle $P(X)$, and so on the space of sections. The fixed point set of Z on $P(X)$ is $P(X)^Z = P(X^Z)$, the associated bundle with fiber the fixed point set $X^Z \subset X$.

Proposition 3.1. *Let G, Z, P, X, C be as above. The action of Z on the space of sections of $P(X)$ preserves Mundet semistability and so induces an action of Z on $\overline{\mathcal{M}}_n^G(C, X)$.*

Proof. The action of Z on P induces an action on parabolic reductions, preserving the associated graded bundles and morphisms and so the Mundet weight of Definition 1.3. $\square$

The following is straight-forward, and similar to the case of stable maps in Kontsevich [39]:

Proposition 3.2. *Let G, Z, P, X, C be as above. The fixed point locus for the action of Z on $\overline{\mathcal{M}}_n^G(C, X)$ is the substack whose objects are pairs $(p : P \rightarrow C, u : \hat{C} \rightarrow P(X))$ such that*

- (a) u takes values in $P(X^Z)$ on the principal component C_0 ;
- (b) for any bubble component $C_i \subset \hat{C}$ mapping to a point in C , u maps C_i to a one-dimensional orbit of Z on $P(X)$; and
- (c) any node of $\hat{C}$ maps to the fixed point set $P(X^Z)$.

3.2. Wall-crossing terms. In this subsection we explain the wall-crossing terms in the main result Theorem 1.10 and state the generalization to arbitrary group actions Theorem 3.6 below. Let X be a smooth projective G -variety equipped with polarizations $L_{\pm} \rightarrow X$.

Definition 3.3 (Fixed Gauged Maps). For each $\zeta \in \mathfrak{g}$, let $\overline{\mathcal{M}}_n^{G_{\zeta}}(C, X, \zeta, d)$ denote the stack of Mundet-semistable morphisms from C to X/G_{ζ} that are $\mathbb{C}_{\zeta}^{\times}$ -fixed and take values in X^{ζ} on the principal component. Via the inclusion $G_{\zeta} \rightarrow G$ the universal curve over $\overline{\mathcal{M}}_n^{G_{\zeta}}(C, X, \zeta, d)$ admits a morphism to X/G , and the tangent complex $T(X/G)$ defines an object $\text{Ind}(T(X/G))$ in the bounded derived category of $\overline{\mathcal{M}}_n^{G_{\zeta}}(C, X, \zeta, d)$ with an action of $\mathbb{C}_{\zeta}^{\times}$. We denote by $\text{Ind}(T(X/G))^+ \subset \text{Ind} T(X/G)$ the moving part of $\text{Ind}(T(X/G))$ with respect to the action of $\mathbb{C}_{\zeta}^{\times}$ and $\epsilon_+(T(X/G)) = \text{Eul}(\text{Ind}(T(X/G))^+)$ its Euler class, as in (7), but in the more general setting considered here.

Remark 3.4. (Large-area limit of fixed gauged maps) The Mundet semistability condition for fixed gauged maps simplifies somewhat in the limit $\rho \rightarrow \infty$. Suppose that $X^{\zeta, t}$ is the component of X^{ζ} that is L_t -semistable. For any class d there exists a ρ_0 such that for $\rho > \rho_0$, any Mundet-semistable fixed map for polarization L_t must consist of a principal component mapping to $X^{\zeta, t}/G_{\zeta}$ and bubbles mapping to X/G_{ζ} . Indeed, Mundet semistability for L^{ρ} implies that the Hilbert weight $\mu_H(\sigma, \lambda)$ is at most ρ^{-1} times minus the Ramanan weight $\mu_R(\sigma, \lambda)$, for any σ, λ . In particular, this holds $\lambda = -\zeta$ for ζ antidominant and σ the trivial parabolic reduction, in which case the Ramanan weight for a pair (P, u) with respect to (σ, λ) is simply $-\langle c_1(P), \zeta \rangle$. The latter is bounded by a constant $c(d)$ depending on $d \in H_2^G(X, \mathbb{Z})$, since $c_1(P)$ is the projection of d onto $H_2(BG)$. So the Hilbert weight $\mu_M(\sigma, \lambda)$ is less than $c(d)\rho^{-1}$. It follows for any $\rho > \rho_0$ where

$$\rho_0 := c(d) \inf_{s \neq t} |\langle \Phi(X^{\zeta, t}), \zeta \rangle - \langle \Phi(X^{\zeta, s}), \zeta \rangle|^{-1},$$

and any L_t -semistable pair (P, u) , the section u takes values in $P(X^{\zeta, t})$. (In algebraic language, this is $c(d)$ times the inverse of the smallest difference in Hilbert-Mumford weights for ζ for different fixed point components.) To summarize, in the large-area limit $\rho \rightarrow \infty$, the set of values of t for which $\overline{\mathcal{M}}_n^{G_\zeta}(C, X, L_t, \zeta, d)$ is non-empty coincides with the set of values of t for which $X^{\zeta, t}$ is non-empty, and the principal component of any object (P, u) of $\overline{\mathcal{M}}_n^{G_\zeta}(C, X, L_t, \zeta, d)$ takes values in $P(X^{\zeta, t})$.

Conversely, suppose that stable=semistable for the G -action on $\mathbb{P}(L_- \oplus L_+)$. Suppose that (P, u) is $G_\zeta/\mathbb{C}_\zeta^\times$ -semistable and u takes values in $P(X^{\zeta, t})$ on the principal component. Then (P, u) is $L_{t'}^\rho$ -semistable for G_ζ for sufficiently large ρ for a unique t' in a small neighborhood of t . Indeed, the stable=semistable assumption implies that Hilbert-Mumford weight for the action of $\mathbb{C}_\zeta^\times$ on $X^{\zeta, t}$ is non-constant (that is, $X^{\zeta, t}$ is semistable for a single value of t , see Remark 2.2). It follows that $C_\zeta^\times$ -semistability of (P, u) for t holds for s close to t and ρ sufficiently large so that $\mu_H(\sigma, \zeta, s) = -\rho^{-1}\mu_R(\sigma, \zeta, s)$, while $G_\zeta/\mathbb{C}_\zeta^\times$ -semistability holds by assumption. In general, semistability for a product of groups is not equivalent to semistability for the factors but this is true if one of the factors acts trivially, as in the present case.

We will construct in the proof below a perfect relative obstruction theory on $\overline{\mathcal{M}}_n^{G_\zeta}(C, X, L_t, \zeta)$. This construction allows the following definition:

Definition 3.5 (Fixed point contributions to wall-crossing for Gromov-Witten invariants). Virtual integration over the stacks $\overline{\mathcal{M}}_n^{G_\zeta}(C, X, L_t, \zeta, d)$, $d \in H_2^G(X, \mathbb{Z})$ defines a “fixed point contribution”

$$(13) \quad \tau_{X, \zeta, t}^{G_\zeta} : QH_{G, \text{fin}}(X) \rightarrow \tilde{\Lambda}_X^G[\xi, \xi^{-1}],$$

$$\alpha \mapsto \sum_{d \in H_2^G(X, \mathbb{Z})} \sum_{n \geq 0} \int_{[\overline{\mathcal{M}}_n^{G_\zeta}(C, X, L_t, \zeta, d)]} (q^d/n!) \text{ev}^*(\alpha, \dots, \alpha) \cup \epsilon_+(T(X/G))^{-1}$$

for $\alpha \in H_G(X)$, extended by linearity over $\Lambda_X^{G, \text{fin}}$, where we omit the restriction map $H_{G, \text{fin}}(X) \rightarrow H_{G_\zeta}(X)$ to simplify notation.

Theorem 3.6 (quantum Kalkman formula, arbitrary group case). *Suppose that X is equipped with polarizations $L_\pm$ so stable=semistable for the action of G on $\mathbb{P}(L_- \oplus L_+)$. Then the Gromov-Witten invariants of $X//_\pm G$ are related by a sum of twisted gauged Gromov-Witten invariants for subgroups $G_\zeta \subset G$*

$$(14) \quad \tau_{X//_+ G} \kappa_{X, +}^G - \tau_{X//_- G} \kappa_{X, -}^G = \lim_{\rho \rightarrow \infty} \sum_{[\zeta], t \in (-1, 1)} \frac{|W_\zeta|}{|W_{\mathbb{C}\zeta}|} \text{Resid}_\xi \tau_{X, \zeta, t}^{G_\zeta},$$

where the sum over $[\zeta]$ is over equivalence classes of unparameterized one-parameter subgroups generated by $\zeta \in \mathfrak{g}$.

We already gave a simple Example 1.11 of this formula in the introduction. We give another Fano example:

Example 3.7. (Quantum powers of the first Chern class for the blow-up of the projective plane) Suppose that $G = (\mathbb{C}^\times)^2$ acts on $X = \mathbb{C}^4$ with weights $(1, 0), (1, 0), (1, 1)(0, 1)$. Consider the path from $(-1, 2)$ to $(2, -1)$ in $H_G^2(X) \cong \mathbb{Q}^2$ crossing through the chambers with git quotients $\emptyset, \mathbb{P}^2, \text{Bl}(\mathbb{P}^2), \emptyset$. We denote by $X//_-G$ resp. $X//_+G$ the second resp. third quotient. The quantum Kirwan morphism for $\mathbb{P}^2, \text{Bl}(\mathbb{P}^2)$ has no quantum corrections, since these varieties are Fano, and so

$$(15) \quad D_0 \kappa_{X, \pm}^G(c_1^G(X)) = c_1(X//_{\pm}G).$$

We consider the wall-crossing formula for invariants with 5 fixed markings, corresponding to the small quantum product $c_1(X//_{\pm}G)^{\star 5} \in QH(X//_{\pm}G)$. By Example 1.5, the moduli space of gauged maps is, after fixing the locations of the markings, the quotient of the space of sections $H^0(\mathbb{P}, \mathcal{O}_C(d) \times_{\mathbb{C}^\times} \mathbb{C}^4)$ by G , with stability condition corresponding to the stability condition for X , see [55]. We take $d = (1, 0)$, so that

$$H^0(\mathbb{P}, \mathcal{O}_C(d) \times_{\mathbb{C}^\times} \mathbb{C}^4) = \mathbb{C}_{(0,1)} \oplus \mathbb{C}_{(1,1)}^{\oplus 2} \oplus \mathbb{C}_{(0,1)}^{\oplus 4}$$

(or $\mathbb{C}^7$ for short) where for any weight μ , $\mathbb{C}_\mu$ denotes the one-dimensional representation with weight μ . The moduli spaces of gauged maps corresponding to the various chambers are therefore the empty set, $\mathbb{P}^3$, its blow-up along a $\mathbb{P}$, and the empty set again. We consider the sequence of polarizations corresponding to the vectors $(-1, 2), (1, 2), (2, 1), (2, -1)$. The first wall-crossing term for degree $(1, 0)$ invariants corresponds to the direction $\zeta = (1, 0)$, for which there is a unique ζ -fixed stable map with normal weights $(1, 0)$ with multiplicity 4 and $(1, 1)$ with multiplicity 2. The wall-crossing term is

$$\text{Resid}_{\xi_1} \frac{(3\xi_1 + 2\xi_2)^5}{\xi_1^4(\xi_1 + \xi_2)^2} = 243.$$

Using (15) this implies that $c_1(\mathbb{P}^2)^{\star 5} \supset 243q_1[\text{pt}]$ which is shorthand for saying that the coefficient of $q_1[\text{pt}]$ is 243; here $\star$ is the small quantum product, as expected since

$$c_1(\mathbb{P}^2)^{\star 5} = (3\omega)^{\star 5} = 243\omega^{\star 3} \star \omega^{\star 2} = 243q_1[\text{pt}].$$

The second wall-crossing term corresponds to the change of quotient from $\mathbb{P}^2$ to $\text{Bl}(\mathbb{P}^2)$ with $\zeta = (1, -1)$ is (after fixing the five points on $\mathbb{P}$) an integration over $\overline{\mathcal{M}}_0^G(\mathbb{P}, X, \zeta, t) \cong \mathbb{P}$, the quotient of the ζ -fixed summand $\mathbb{C}_{(1,1)}^{\otimes 2}$ by $\mathbb{C}^\times$,

$$\frac{1}{2} \text{Resid}_\xi \int_{[\mathbb{P}]} \kappa_{\mathbb{C}^7}^{\mathbb{C}^\times} \frac{(3\xi_1 + 2\xi_2)^5}{\xi_1^4 \xi_2}$$

where $\kappa_{\mathbb{C}^7}^{\mathbb{C}^\times}$ is the descent map in equivariant cohomology from $\mathbb{C}^7$. Using that $\kappa_{\mathbb{C}^6, \mathbb{C}^\times}(\xi_1)$ is the generator ω of $H^2(\mathbb{P})$ this gives

$$\begin{aligned} \frac{1}{2} \text{Resid}_\xi \int_{[\mathbb{P}]} \frac{(5\omega + \xi)^5}{(\omega + \xi)^4(\omega - \xi)} &= \text{Resid}_\xi \int_{[\mathbb{P}]} -\xi^{-5}(25\omega\xi^4 + \xi^5)(1 - 4\omega/\xi)(1 + \omega/\xi) \\ &= 11. \end{aligned}$$

It follows that $c_1(\mathrm{Bl}(\mathbb{P}^2))^{\star 5} \supset 232q_1[\mathrm{pt}]$. The last wall-crossing term for the degree $(1, 0)$ invariant is

$$\mathrm{Resid}_\xi \int_{[\mathbb{P}^3]} \kappa_{\mathbb{C}^\times}^7 \frac{(3\xi_1 + 2\xi_2)^5}{\xi_2(\xi_1 + \xi_2)^2} = 232.$$

This is as expected since the quotient is empty in the last chamber. One can verify that the expansion of $c_1(\mathrm{Bl}(\mathbb{P}^2))^{\star 5}$ contains $232q_1[\mathrm{pt}]$ using the known quantum multiplication table for $\mathrm{Bl}(\mathbb{P}^2)$ from Crauder-Miranda [16]:

$\star$	e	f	p
e	$-p + eq^e + xq^f$	$p - eq^e$	$f q^f$
f		eq^e	xq^{e+f}
p			$(e + f)q^{e+f}$

where e, f, p, x are the exceptional resp. fiber resp. point resp. fundamental classes respectively, $q_1 = q^{e+f}$, and using a little help from Mathematica. (We thank Eric Malm for teaching us how to get Mathematica to compute these coefficients.)

3.3. Construction of a master space. By the adiabatic limit theorem 1.8, to prove the wall-crossing formula 3.6 it suffices to prove a formula for the difference of gauged potentials. The following result is a algebro-geometric generalization of a wall-crossing formula of Cieliebak-Salamon [12] for gauged Gromov-Witten invariants of quotients of vector spaces defined using symplectic geometry. We will deduce our main result Theorem 3.6 by taking the large area limit $\rho \rightarrow \infty$ of the following Theorem:

Theorem 3.8 (Wall-crossing for gauged Gromov-Witten potentials). *Let X be a smooth projective G -variety. Suppose that $L_\pm \rightarrow X$ are polarizations such that semistable=stable for the stack $\overline{\mathcal{M}}_n^G(C, X, L_-, L_+)$ of 3.9. Then the gauged Gromov-Witten potentials are related by*

$$(16) \quad \tau_{X,+}^G - \tau_{X,-}^G = \sum_{[\zeta], t \in (-1, 1)} \frac{|W_\zeta|}{|W_{\mathbb{C}\zeta}|} \mathrm{Resid}_\xi \tau_{X, \zeta, t}^{G_\zeta}$$

where the sum is over equivalence classes $[\zeta]$ of unparametrized one-parameter subgroups generated by $\zeta \in \mathfrak{g}$, and the same Mundet semistability parameter should be used to define the potentials on both sides of the equation.

The proof of the wall-crossing formula 3.8 depends on the construction of *master space* in the sense of 2.1 whose quotients are the moduli spaces of Mundet stable gauged maps.

Proposition 3.9 (Existence of a master space). *Under suitable stable=semistable conditions, there exists a proper Deligne-Mumford $\mathbb{C}^\times$ -stack with perfect relative obstruction theory $\overline{\mathcal{M}}_n^G(C, X, L_+, L_-, d)$ whose $\mathbb{C}^\times$ -fixed substack is the union of $\overline{\mathcal{M}}_n^G(C, X, L_-, d)$, $\overline{\mathcal{M}}_n^G(C, X, L_+, d)$ and the images of the stacks $\overline{\mathcal{M}}_n^{G_\zeta}(C, X, L_t, \zeta, d)$ for $t \in (-1, 1)$ of Definition 3.5. Furthermore, $\overline{\mathcal{M}}_n^G(C, X, L_+, L_-, d)$ admits an embedding in a non-singular Deligne-Mumford stack.*

The proof will be given after several constructions. First recall the quot-scheme compactification of Mundet semistable morphisms from C to X/G constructed by Schmitt [50]:

Definition 3.10 (Quot-scheme compactification of gauged maps). In the case $X = \mathbb{P}^{n-1}$ and $G = GL(n)$, a *projective bump* over a smooth projective curve C is a datum consisting of a vector bundle $E \rightarrow C$; a line bundle $L \rightarrow C$; and a surjective morphism $E \rightarrow L$. More generally, if G is an arbitrary group embedded in $GL(r)$, and X is a smooth projective G -variety embedded in $\mathbb{P}^{r-1}$ then a projective bump is a projective $GL(r)$ -bump and a reduction of the frame bundle $\text{Fr}(E^\vee)$ to G such that the corresponding injection $L^\vee \rightarrow E^\vee$ takes values in the associated X -bundle $\text{Fr}(E^\vee) \times_G X \subset \text{Fr}(E^\vee) \times_G \mathbb{P}^{r-1}$. A bump is *Mundet semistable* if it satisfies the inequality of Definition 1.3 for every pair (σ, λ) consisting of a parabolic reduction σ and antidominant coweight λ .

Let $\overline{\mathcal{M}}^{G, \text{quot}}(C, X, d)$ denote the stack of Mundet semistable projective bumps of class $d \in H_2^G(X, \mathbb{Z})$ over C with target X .

Example 3.11. In the toric case discussed in Example 1.5 there are no non-constant holomorphic maps from curves to X and so $\overline{\mathcal{M}}^{G, \text{quot}}(C, X, d) = \mathcal{M}_0^G(C, X, d) = H^0(C, \mathcal{O}_C(d) \times_G X) // G$. That is, the moduli space of maps from irreducible domain is already compact and does not require compactification.

The following summarizes results of Schmitt [50]:

Proposition 3.12 (git construction of projective bumps). [50] $\overline{\mathcal{M}}^{G, \text{quot}}(C, X, d)$ admits a presentation as a git quotient of a framed moduli space $\overline{\mathcal{M}}^{G, \text{quot}, \text{fr}}(C, X, d)$ (consisting of bundles that can be written as quotients of a given trivial bundle on C , which we omit from the notation) by the action of a group H acting on the space of framings, with respect to a polarization $D(L)$ on $\overline{\mathcal{M}}^{G, \text{quot}, \text{fr}}(C, X, d)$ depending on the choice of polarization $L \rightarrow X$. If stable=semistable, then $\overline{\mathcal{M}}^{G, \text{quot}}(C, X, d)$ is a proper Deligne-Mumford stack with a projective coarse moduli space.

The stack $\overline{\mathcal{M}}^{G, \text{quot}}(C, X, d)$ seems not to be virtually smooth in general and so, it seems, cannot be expected to admit a good virtual fundamental class. We denote by $\overline{\mathfrak{M}}^{G, \text{quot}}(C, X, d)$ the stack-theoretic quotient, $\overline{\mathfrak{M}}^{G, \text{quot}}(C, X, d) := \overline{\mathcal{M}}^{G, \text{quot}, \text{fr}}(C, X, d) / H$ which contains $\overline{\mathcal{M}}^{G, \text{quot}}(C, X, d)$ as an open substack. In order to study the dependence on polarization, suppose we are given two polarizations $L_\pm \rightarrow X$ and denote by $\mathcal{O}_{\mathbb{P}(D(L_-) \oplus D(L_+))}(1) \rightarrow \mathbb{P}(D(L_-) \oplus D(L_+))$ the relative hyperplane bundle as in (10). Consider the git quotient

$$\overline{\mathcal{M}}^{G, \text{quot}}(C, X, L_+, L_-, d) := \mathbb{P}(D(L_-) \oplus D(L_+)) // H.$$

We call an object of $\overline{\mathcal{M}}^{G, \text{quot}}(C, X, d, L_-, L_+)$ a *polarized bump*, since generically an object of it is equivalent to a bump together with a point in the line bundle over it. From the geometric invariant theory construction, it follows:

Proposition 3.13 (Stack of polarized bumps). *If stable=semistable then the stack $\overline{\mathcal{M}}^{G,\text{quot}}(C, X, d, L_-, L_+)$ is a smooth proper Deligne-Mumford stack with a projective coarse moduli space.*

We denote by $\overline{\mathcal{M}}^{G,\text{quot}}(C, X, d, L_t) \subset \overline{\mathfrak{M}}^{G,\text{quot}}(C, X, d)$ the stable locus of the bundle $D(L_t)$ for $L_t = L_-^{(1-t)/2} L_+^{(1+t)/2}$ depending on $t \in [-1, 1]$, and by

$$\overline{\mathfrak{M}}^{G,\text{quot}}(C, X, L_+, L_-, d) := \mathbb{P}(D(L_-) \oplus D(L_+))/H$$

the Artin quotient stack. By Lemma 2.1

Proposition 3.14 (Master space for bumps). *The residual action of $\mathbb{C}^\times$ on the stack $\overline{\mathcal{M}}^{G,\text{quot}}(C, X, d, L_-, L_+)$ has git quotient with respect to the polarization $D(L_t)$ isomorphic to $\overline{\mathcal{M}}^{G,\text{quot}}(C, X, d, L_t)$.*

We wish to obtain similar results for the Kontsevich-style compactification, which is the one having a virtual fundamental class with the expected properties. The projection

$$(17) \quad \pi : \overline{\mathfrak{M}}^{G,\text{quot}}(C, X, L_+, L_-) \rightarrow \overline{\mathfrak{M}}^{G,\text{quot}}(C, X)$$

is a $\mathbb{P}$ -bundle over $\overline{\mathfrak{M}}^{G,\text{quot}}(C, X)$. Let $\overline{\mathfrak{M}}_n^G(C, X, [-1, 1]) \subset \overline{\mathfrak{M}}_n^G(C, X)$ denote the Artin stack of gauged maps that are semistable with respect to the stability condition defined by L_t for some $t \in [-1, 1]$:

$$\overline{\mathfrak{M}}_n^G(C, X, [-1, 1]) := \bigcup_{t \in [-1, 1]} \overline{\mathcal{M}}_n^G(C, X, L_t).$$

Lemma 3.15. *There is a relative Givental morphism from $\overline{\mathfrak{M}}_n^G(C, X, [-1, 1])$ to $\overline{\mathfrak{M}}^{G,\text{quot}}(C, X)$ extending the forgetful morphism on the open stratum (if non-empty).*

Proof. The morphism, roughly speaking, replaces each bubble tree which maps to a point in C with a base point for the linear system corresponding to the quotient. The relative Givental morphism is constructed more generally in Popa-Roth [48]. Each object in $\overline{\mathfrak{M}}_n^G(C, X, [-1, 1])$ is a pair of a bundle $P \rightarrow C \times S$ with stable section $u : \hat{C} \rightarrow P(X)$, where $\hat{C} \rightarrow S$ is a nodal curve; the Givental morphism replaces u with the corresponding quotient. $\square$

Remark 3.16. There is no morphism from $\overline{\mathfrak{M}}_n^G(C, X, d)$ to $\overline{\mathfrak{M}}^{G,\text{quot}}(C, X, d)$ since the domain has objects with infinitely many isomorphism classes of bundles, while the isomorphism classes of bundles in the target form a finite set.

Definition 3.17 (Polarized gauged maps). A *Mundet semi-stable polarized gauged map* is an object in the fiber product of the relative Givental morphism with the projection from the master space for projective bumps:

$$\overline{\mathcal{M}}_n^G(C, X, L_+, L_-) := \overline{\mathfrak{M}}_n^G(C, X, [-1, 1]) \times_{\overline{\mathfrak{M}}^{G,\text{quot}}(C, X)} \overline{\mathcal{M}}^{G,\text{quot}}(C, X, L_+, L_-)$$

that is, a polarized gauged bump together with a lift of the underlying bump to the moduli stack of stable gauged maps that are Mundet-semistable for some $t \in [-1, 1]$.

Proposition 3.18. *If stable=semistable for polarized gauged maps, then for any $d \in H_2^G(X, \mathbb{Z})$, $\overline{\mathcal{M}}_n^G(C, X, L_+, L_-, d)$ is a smooth proper Deligne-Mumford stack.*

Proof. Since $\overline{\mathcal{M}}_n^G(C, X, L_+, L_-, d) \rightarrow \overline{\mathcal{M}}^{G, \text{quot}}(C, X, L_+, L_-, d)$ is proper and Deligne-Mumford, and the codomain is proper Deligne-Mumford under the stable=semistable assumption. Smoothness is immediate from the fiber product description, since the second morphism is an open embedding. $\square$

3.4. Analysis of the fixed point contributions. In this subsection, we deduce the quantum wall-crossing formula by applying virtual localization to the master space $\overline{\mathcal{M}}_n^G(C, X, L_+, L_-, d)$ constructed in the previous subsection. Under the given hypotheses the stack $\overline{\mathcal{M}}_n^G(C, X, L_+, L_-, d)$ is a proper Deligne-Mumford stack and an open substack of Artin stack $\overline{\mathfrak{M}}_n^G(C, X, L_+, L_-, d)$ defined by dropping the semistability condition.

Proposition 3.19 (Obstruction theory for polarized gauged maps). *For any $d \in H_2^G(X, \mathbb{Z})$, the Artin stack $\overline{\mathfrak{M}}_n^G(C, X, L_+, L_-, d)$ has a relative obstruction theory over the moduli stack $\overline{\mathfrak{M}}_n(C)$ of parametrized prestable curves. If stable=semistable then the restriction of this obstruction theory to $\overline{\mathcal{M}}_n^G(C, X, L_+, L_-, d)$ is perfect.*

Proof. The complex in the relative obstruction theory is denoted $?$ in the diagram below, constructed from the following commutative diagram of complexes of coherent sheaves. We use $T^\vee$ resp. $T_\pi^\vee$ for the cotangent complex resp. relative cotangent complex for the morphism (17) to avoid conflict with previous notation.

$$\begin{array}{ccccccc}
 T_\pi^\vee & \longleftarrow & ? & \longleftarrow & \pi^* R p_* e^* T(X/G) & \longleftarrow & T_\pi^\vee[1] \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 T_\pi^\vee & \longleftarrow & T_{\overline{\mathfrak{M}}_n^G(C, X, L_+, L_+)}^\vee & \longleftarrow & \pi^* T_{\overline{\mathfrak{M}}_n^G(C, X, [-1, 1])}^\vee & \longleftarrow & T_\pi^\vee[1]
 \end{array}$$

where the horizontal lines are exact triangles and the second vertical arrow $\hat{\phi}$ exists by the axioms of a triangulated category. The map $\hat{\phi}$ satisfies the axioms of an obstruction theory by the five lemma applied to the cohomology of the above diagram. If the automorphisms are finite then the relative obstruction theory is perfect. Indeed for the base H^1 is identified with the dual of the Lie algebra of the group of automorphisms [55, Example 6.7(c)], and this carries over to the total space by the long exact sequence. $\square$

Lemma 3.20. *For any $d \in H_2^G(X, \mathbb{Z})$, $\overline{\mathcal{M}}_n^G(C, X, L_+, L_-, d)$ admits an embedding in a non-singular Deligne-Mumford stack.*

Proof. Consider an embedding $G \rightarrow GL(k)$ and G -equivariant embedding $X \rightarrow \mathbb{P}^{k-1}$ for some k . Let $\overline{\mathcal{U}}_n^{G, \text{quot}, \text{fr}}(C, X, L_+, L_-, d)$ denote the universal bundle over the moduli space of framed bumps $\overline{\mathcal{M}}_n^{G, \text{quot}, \text{fr}}(C, X, L_+, L_-, d)$, equipped with the action of H by

changing the framings as in Proposition 3.12. Then $\overline{U}_n^{G, \text{quot}, \text{fr}}(C, X, L_+, L_-, d) \times_G X$ embeds in the projectivization $\overline{U}_n^{GL(k), \text{quot}, \text{fr}}(C, X, L_+, L_-, d) \times_{GL(k)} \mathbb{P}^{k-1}$. The latter is projective by Schmitt's construction (it is the pull-back of the universal bundle on quot scheme) and so embeds H -equivariantly in some $\mathbb{P}^N$. Then $\overline{\mathcal{M}}_n^G(C, X, L_+, L_-, d)$ is an embedded substack of $\overline{\mathcal{M}}_{g,n}(\mathbb{P}^N)/H$, with objects given by stable maps that are compositions of stable sections of $\overline{U}_n^{G, \text{quot}, \text{fr}}(C, X, L_+, L_-, d) \times_G X/H$ with the inclusion into $\mathbb{P}^N/H$. Since $\overline{\mathcal{M}}_{g,n}(\mathbb{P}^N)$ is a non-singular Deligne-Mumford stack, the quotient $\overline{\mathcal{M}}_{g,n}(\mathbb{P}^N)/H$ is a non-singular Artin stack. The group H acts locally freely on an open subset $\overline{\mathcal{M}}_{g,n}(\mathbb{P}^N)^{\text{reg}} \subset \overline{\mathcal{M}}_{g,n}(\mathbb{P}^N)$ by the stable=semistable assumption and the quotient $\overline{\mathcal{M}}_{g,n}(\mathbb{P}^N)^{\text{reg}}/H$ is Deligne-Mumford. $\square$

Proposition 3.21 (Wall-crossing for gauged Gromov-Witten invariants, preliminary version). *Let X be a smooth projective G -variety equipped with polarizations $L_{\pm} \rightarrow X$. Suppose that stable=semistable for the master space $\overline{\mathcal{M}}_n^G(C, X, L_+, L_-, d)$. Then for any $\alpha \in H_G(X)^n$,*

$$(18) \quad \int_{[\overline{\mathcal{M}}_n^G(C, X, L_+, d)]} \text{ev}^* \alpha - \int_{[\overline{\mathcal{M}}_n^G(C, X, L_-, d)]} \text{ev}^* \alpha \\ = \text{Resid}_{\xi} \left(\sum_F \int_{[F]} \iota_F^* \text{ev}^* \alpha \cup \text{Eul}_{\mathbb{C}^\times}(\nu_F)^{-1} \right)$$

where F ranges over the fixed point components of $\mathbb{C}^\times$ on $\overline{\mathcal{M}}_n^G(C, X, L_+, L_-, d)$ not equal to $\overline{\mathcal{M}}_n^G(C, X, L_{\pm}, d)$ with normal complex ν_F .

Proof. By virtual localization applied to the $\mathbb{C}^\times$ -action on $\overline{\mathcal{M}}_n^G(C, X, L_+, L_-, d)$, as in the Kiem-Li fixed point formula Theorem 2.6. $\square$

Next we describe the moduli spaces of $\mathbb{C}^\times$ -fixed gauged maps in terms of gauged maps with smaller structure group. Let $\overline{\mathcal{M}}_n^{G_\zeta}(C, X, L_t, \zeta)$ denote the category of ζ -fixed gauged maps whose principal component is semistable for the polarization given by L_t for some $t \in (-1, 1)$, as in Definition 3.3.

Proposition 3.22 (Fixed points as reducible gauged maps). *Any $\mathbb{C}^\times$ -fixed component of $\overline{\mathcal{M}}_n^G(C, X, L_+, L_-)$ is in the image of $\overline{\mathcal{M}}_n^{G_\zeta}(C, X, L_t, \zeta)$ in $\overline{\mathcal{M}}_n^G(C, X, L_+, L_-)$ for some $t \in (-1, 1)$ where $\zeta \in \mathfrak{g}$ is a non-zero element, G_ζ is stabilizer, and $\mathbb{C}_\zeta^\times \subset G_\zeta$ the unparametrized one-parameter subgroup generated by ζ , consisting of maps $u : \hat{C} \rightarrow X/G_\zeta$ taking values in X^ζ/G_ζ on the principal component, and X/G_ζ on the bubbles. Furthermore, for $\zeta \in \mathfrak{t}$ the map $\overline{\mathcal{M}}_n^{G_\zeta}(C, X, L_t, \zeta) \rightarrow \overline{\mathcal{M}}_n^G(C, X, L_+, L_-)$ has fiber isomorphic to $W_{\mathbb{C}_\zeta^\times}/W_\zeta$.*

Proof. Any fixed point of $\mathbb{C}^\times$ on $\overline{\mathcal{M}}_n^G(C, X, L_+, L_-)$ is necessarily in the fiber over a gauged map $\pi : P \rightarrow C, u : \hat{C} \rightarrow P(X)$ with infinite automorphism group. The infinitesimal automorphism generated by the action of $\mathbb{C}^\times$ is a section of the adjoint

bundle $P(\mathfrak{g})$, given by an element $\zeta \in P(\mathfrak{g})_z \cong \mathfrak{g}$ at a base point $z \in C$. The structure group of P reduces to the centralizer G_ζ , and being fixed by the automorphism the section u takes values in the fixed point set $P(X^\zeta) = P(X)^\zeta$ of ζ on the principal component. The fiber of the morphism is $N(G_\zeta)_{\mathbb{C}_\zeta^\times}/G_\zeta = W_{\mathbb{C}_\zeta^\times}/W_\zeta$ for $\zeta \in \mathfrak{t}$, corresponding to pairs (P, u) related by G -isomorphisms that are not G_ζ -isomorphisms. $\square$

Remark 3.23. We may describe the fixed point locus as a stack of “bubble trees” as follows: There is an isomorphism

$$\overline{\mathcal{M}}_n^{G_\zeta}(C, X, L_t, \zeta) \rightarrow \bigcup_{r, [I_1, \dots, I_r]} \left(\mathcal{M}_r^{G_\zeta, \text{fr}}(C, X^{\zeta, t}) \times_{(X^\zeta)^r} \prod_{j=1}^r \overline{\mathcal{M}}_{|I_j|+1}(X) \right)^\zeta / (G_\zeta)^r$$

where the right-hand-side denotes the fixed point set of the action $\mathbb{C}_\zeta^\times$ on the moduli stack of gauged maps to $X^{\zeta, t}$ with bubbles in X . Indeed, by definition each object of $\overline{\mathcal{M}}_n^{G_\zeta}(C, X, L_t, \zeta)$ consists of a principal component mapping to $X^{\zeta, t}/G_\zeta$ and a collection of bubble trees in X fixed (up to reparametrization) by the action of $\mathbb{C}_\zeta^\times$.

Corollary 3.24. (Obstruction theory for the fixed point components) $\overline{\mathcal{M}}_n^{G_\zeta}(C, X, L_t, \zeta)$ is an Artin stack, and if every automorphism group is finite modulo $\mathbb{C}_\zeta^\times$, each substack with fixed homology class $d \in H_2^{G_\zeta}(X, \mathbb{Z})$ is a proper Deligne-Mumford stack with a relatively perfect obstruction theory over $\overline{\mathcal{M}}_n(C) \times B\mathbb{C}_\zeta^\times$ whose complex is $\text{Ind}(T(X/G)_+)$.

Proof. The relatively perfect obstruction theory $\overline{\mathcal{M}}_n^{G_\zeta}(C, X, L_t, \zeta)$ is pulled back from that on the $\mathbb{C}^\times$ -fixed point set in $\overline{\mathcal{M}}_n^G(C, X, L_-, L_+)$ in Proposition 3.22, which exists by the construction of relatively perfect obstruction theories on fixed point loci in [27]. $\square$

Let $[\overline{\mathcal{M}}_n^{G_\zeta}(C, X, L_t, \zeta, d)]$ denote the virtual fundamental class in the homology of the coarse moduli space resulting from Corollary 3.24. Integration with respect to these classes yields ζ -fixed gauged Gromov-Witten invariants of Definition 3.5. These gauged Gromov-Witten invariants satisfy splitting properties similar to the gauged Gromov-Witten invariants defined by integration over $\overline{\mathcal{M}}_n^G(C, X)$ constructed in [25], [55], that we will not discuss here. The ζ -fixed gauged Gromov-Witten invariants that appear in the wall-crossing formula involve further twists by Euler classes of index bundles $(Rp_*e^*T(X/G))^+$, that is, the moving part of $Rp_*e^*T(X/G)$. This completes the construction of the fixed point potential in Definition 3.5. We remark that the resulting fixed point potential $\tau_{X, \zeta, t}^{G_\zeta}$ takes values in $\tilde{\Lambda}_X^G$ rather than in Λ_{X, L_t}^G because Gromov compactness fails for gauged maps in the case that a central subgroup $\mathbb{C}_\zeta^\times$ acts trivially. Indeed, in this case, the energy $\langle d, c_1(L) \rangle$ of a gauged map of class d does not determine the isomorphism class of the bundle, since twisting by a character of $\mathbb{C}_\zeta^\times$ does not change the energy.

Theorem 3.6 now follows from Theorem 3.21 and Proposition 3.22, and the observation that stable=semistable for X implies stable=semistable for gauged maps in the limit $\rho \rightarrow \infty$, discussed in [55] and, for fixed gauged maps, in Remark 3.4.

Remark 3.25. The right-hand-side of Theorem 3.6 can also be re-written using the quantum Kirwan map for $QH_{G_\zeta/\mathbb{C}^\times}(X^{\zeta,t}) \rightarrow QH(X^{\zeta,t} // (G_\zeta/\mathbb{C}^\times))$, using the adiabatic limit theorem for $(G_\zeta/\mathbb{C}^\times)$ -gauged maps. However, in our examples the gauged Gromov-Witten invariants are always easier to compute than the Gromov-Witten invariants of the git quotients, so we have left the formula as written.

4. INVARIANCE UNDER CREPANT TRANSFORMATIONS

Ruan and others, see [13], conjectured that *crepant resolutions* induce equivalences in Gromov-Witten theory. More generally, suppose that $Y_\pm$ are smooth proper Deligne-Mumford stacks with projective coarse moduli spaces related by a birational equivalence ϕ . Such a birational equivalence is called *crepant* (or a *K-equivalence*) if ϕ factors into morphisms $\phi_\pm : Z \rightarrow Y_\pm$ from a smooth stack Z with projective coarse moduli space such that the pullbacks of the canonical divisors to Z are equal, as in Kawamata [35]. It is conjectured by Li-Ruan [43], Coates-Ruan [13] and others (perhaps motivated by physics papers such as Witten [54]) that in such a situation (not necessarily arising from geometric invariant theory) the Gromov-Witten theories of Y_- and Y_+ are equivalent, in a sense to be made precise. Many special cases have been proved, see for example Iwao-Lee-Lin-Wang [32], Lee-Lin-Wang [40], Boissière-Mann-Perroni [5], Bryan-Gholampour [9], [8], Bryan-Graber-Pandharipande [11], Coates-Corti-Iritani-Tseng [14] and Coates-Iritani-Tseng [15].

We prove a version Theorem 1.12 of Ruan's conjecture for crepant birational equivalences induced by variation of git. Suppose that X is a smooth projective G -variety, and $X//_\pm G$ are git quotients obtained from polarizations $L_\pm \rightarrow X$. Suppose that stable=semistable for $\mathbb{P}(L_- \oplus L_+)$ so that the master space $\tilde{X} = \mathbb{P}(L_- \oplus L_+) // G$ is a smooth proper Deligne-Mumford stack.

Definition 4.1. A birational transformation of git type $\phi : X//_-G \rightarrow X//_+G$ will be called *crepant* if the sum of the weights of $\mathbb{C}^\times$ on NF , counted with multiplicity, vanishes. More generally, in the case of twisted Gromov-Witten theory associated to a G -vector bundle $E \rightarrow X$, we say that $\phi : X//_-G \rightarrow X//_+G$ is crepant resp. weakly crepant if the same holds for $NF \oplus \text{Ind}(E)$ where $\text{Ind}(E)$ is the index of E as in (5).

It is easy to check that the definition of crepant transformation of git type is a special cases of the definition of crepant transformation (*K-equivalence*) in Kawamata [35] etc. Indeed, Kempf's descent lemma [19, Theorem 2.3] and the crepant condition together imply that the canonical bundle (possibly twisted by the top exterior power of $\text{Ind}(E)$) descends to each singular quotient, from which the canonical bundles on $X//_\pm G$ are pulled back.

The proof of invariance follows from a symmetry of the fixed point contributions under an action of the Picard stack $\text{Pic}(C) := \text{Hom}(C, B\mathbb{C}^\times)$ of line bundles on C defined as follows. The Lie algebra $\mathfrak{g}_\zeta$ has a distinguished factor generated by ζ , and

using an invariant metric the weight lattice of $\mathfrak{g}_\zeta$ has a distinguished factor $\mathbb{Z}$ given by its intersection with the Lie algebra of $\mathbb{C}_\zeta^\times$. After passing to a finite cover, we may assume that $G_\zeta \cong G_\zeta/\mathbb{C}_\zeta^\times \times \mathbb{C}_\zeta^\times$.

Lemma 4.2. (Action of the Picard stack on fixed loci) *In the limit $\rho \rightarrow \infty$ there is a canonical action of $\text{Pic}(C)$ on the moduli stack $\overline{\mathcal{M}}_n^{G_\zeta}(C, X, L_t, \zeta)$ for any $t \in (-1, 1)$. The isomorphism preserves the relative obstruction theories and so the Behrend-Fantechi virtual fundamental classes. Furthermore, the isomorphism preserves the class $\text{ev}^* \alpha$ for any $\alpha \in H_{G_\zeta}(X)^n$.*

Proof. An object of $\overline{\mathcal{M}}_n^{G_\zeta}(C, X, L_t, \zeta)$ consists of a tuple $(P, \hat{C}, u)$ where $P \rightarrow C$ is a G -bundle and $u : \hat{C} \rightarrow P(X)$ is ζ -fixed, in particular, the restriction of u to the principal component of C maps into the fixed point locus X^ζ . Let $\mathbb{C}_\zeta^\times \subset G_\zeta$ denote the subgroup of G_ζ generated by $\zeta \in \mathfrak{g}_\zeta$. Then $\text{Pic}(C)$ acts on $\overline{\mathcal{M}}_n^{G_\zeta}(C, X, L_t, \zeta)$ by

$$Q(P, \hat{C}, u) := (P \times_{\mathbb{C}_\zeta^\times} Q, \hat{C}, u)$$

where we use that $(P \times_{\mathbb{C}_\zeta^\times} Q)(X^\zeta) \cong P(X^\zeta)$ since the action of $\mathbb{C}_\zeta^\times$ on X^ζ is trivial. Because the Mundet weights $\mu_M(\sigma, \lambda)/\rho$ approach $\mu_H(\sigma, \lambda)$ as $\rho \rightarrow \infty$, the limiting Mundet is unchanged by the shift by Q in the limit $\rho \rightarrow \infty$ and so Mundet semistability is preserved, see Remark 3.4. The action of an object Q of $\text{Pic}(C)$ induces an isomorphism

$$(19) \quad \mathcal{S} = \mathcal{S}(Q) : \overline{\mathcal{M}}_n^{G_\zeta}(C, X, L_t, \zeta, d) \rightarrow \overline{\mathcal{M}}_n^{G_\zeta}(C, X, L_t, \zeta, d + \delta)$$

where $\delta = c_1(Q)$. The action of $\text{Pic}(C)$ lifts to the universal curves, denoted by the same notation. Since the obstruction theory on $\overline{\mathcal{M}}_n^{G_\zeta}(C, X, L_t, \zeta, d)$ is the ζ -invariant part of $\text{Ind}(T(X/G))$, the isomorphism preserves the relative obstruction theories and so the Behrend-Fantechi virtual fundamental classes. Since the evaluation map is unchanged by pull-back by $\mathcal{S}$ (up to isomorphism given by twisting by Q), the class $\text{ev}^* \alpha$ is preserved. $\square$

Remark 4.3. In preparation for the proof of Theorem 1.12 we briefly recall a few items from the Schwartz theory of *distributions* for which the standard reference is Hörmander [30]. Recall that if S is the circle then we denote by $\mathcal{D}'(S)$ the space of continuous linear functionals on the smooth functions on S , and by $\mathcal{E}'(S) \subset \mathcal{D}'(S)$ the space of *tempered distributions*. Fourier transform defines an isomorphism of $\mathcal{E}'(S)$ with the space of functions on $\mathbb{Z}$ with polynomial growth. We view q as a coordinate on the punctured plane $\mathbb{C}^\times$. Any formal power series in q, q^{-1} defines a distribution on S , which is tempered if the coefficient of q^d has polynomial growth in d . In particular $\sum_{d \in \mathbb{Z}} q^d$ is the delta function at $q = 1$, and has Fourier transform the constant function with value 1. Any distribution of the form $\sum_{d \in \mathbb{Z}} f(d)q^d$, for $f(d)$ polynomial, is a sum of derivatives of the delta function (since Fourier transform takes multiplication to differentiation) and so is almost everywhere zero.

Proof of Theorem 1.12. We study the dependence of the fixed point contributions $\tau_{X, \zeta, d, t}$ with respect to the Picard action defined in Lemma 4.2. Suppose that Q

is a $\mathbb{C}_\zeta^\times$ -bundle of first Chern class the generator of $H^2(C)$, after the identification $\mathbb{C}_\zeta^\times \rightarrow \mathbb{C}^\times$. We denote the corresponding class in $H_2^{G_\zeta}(X^\zeta)$ by δ . Consider the action of $\mathbb{Z}$ -subgroup of $\text{Pic}(C)$ generated by Q . The contribution of any component $\overline{\mathcal{M}}_n^{G_\zeta}(C, X, L_t, \zeta, d)$ of class $d \in H_2^G(X)$ differs from that from the component induced by acting by $Q^{\otimes r}$, of class $d + r\delta$, by the ratio of Euler classes

$$(20) \quad \text{Eul}_{\mathbb{C}_\zeta^\times}(Rp_*e^*T(X/G)^+) \text{Eul}_{\mathbb{C}_\zeta^\times}(\mathcal{S}^{r,*}Rp_*e^*T(X/G)^+)^{-1}$$

which we now compute. Let $X^{\zeta,t}$ be the component of the fixed point set X^ζ which is semistable for $t \in (-1, 1)$. For simplicity, we assume that $X^{\zeta,t}$ is connected; in general, one should repeat the following argument for each connected component. Consider the decomposition into $\mathbb{C}^\times$ -bundles

$$\nu_{X^{\zeta,t}} = \bigoplus_{i=1}^{m_t} \nu_{X^{\zeta,t},i}$$

where $\mathbb{C}^\times$ acts on $\nu_{X^{\zeta,t},i}$ with non-zero weight $\mu_i \in \mathbb{Z}$. Then $e^*T(X/G)$ is canonically isomorphic to $\mathcal{S}^r e^*T(X/G)$ on the bubble components, since the G -bundles are trivial on those components. Because the pull-back complexes are isomorphic on the bubble components, the difference $(e^*T(X/G))^+ - \mathcal{S}^{r,*}(e^*T(X/G))^+$ is the pullback of the difference of the restrictions to the principal part of the universal curve, that is, the projection on the second factor

$$p_0 : C \times \overline{\mathcal{M}}_n^{G_\zeta}(C, X, L_t, \zeta) \rightarrow \overline{\mathcal{M}}_n^{G_\zeta}(C, X, L_t, \zeta).$$

These restrictions are given by

$$(e^*T(X/G))^{+, \text{prin}} \cong \bigoplus_{i=1}^{m_t} e^* \nu_{X^{\zeta,t},i}, \quad \mathcal{S}^{r,*}(e^*T(X/G))^{+, \text{prin}} \cong \bigoplus_{i=1}^{m_t} e^* \nu_{X^{\zeta,t},i} \otimes (e_C^* Q \times_{\mathbb{C}_\zeta^\times} \mathbb{C}_{r\mu_i})$$

where e_C is the map from the universal curve to C . The projection p_0 is a representable morphism of stacks given as global quotients [2] in which case by Grothendieck-Riemann-Roch [53], [21]

$$(21) \quad \text{Ch}(\mathcal{S}^{r,*} \text{Ind}(T(X/G))^+) = \text{Ch} \left(\text{Ind}(T(X/G))^+ \oplus \bigoplus_{i=1}^{m_t} (z^* e^* \nu_{X^{\zeta,t},i})^{\oplus r\mu_i} \right)$$

where $z : \overline{\mathcal{M}}_n^{G_\zeta}(C, X, L_t, \zeta) \rightarrow C \times \overline{\mathcal{M}}_n^{G_\zeta}(C, X, L_t, \zeta)$ is a constant section of p_0 . The equality of Chern characters above implies an isomorphism in rational K -theory; by the splitting principle we may assume that the $\nu_{X^{\zeta,t},i}$ are line bundles. The difference in Euler classes (20) is therefore given by the Euler class of the last summand in (21)

$$\begin{aligned} \text{Eul}_{\mathbb{C}_\zeta^\times} \left(\bigoplus_{i=1}^{m_t} (z^* e^* \nu_{X^{\zeta,t},i})^{\oplus \mu_i r} \right) &= \prod_{i=1}^{m_t} (\mu_i \xi + c_1(\nu_{X^{\zeta,t},i}))^{\mu_i r} \\ &= \prod_{i=1}^{m_t} (\xi + c_1(\nu_{X^{\zeta,t},i})/\mu_i)^{\mu_i r} \prod_{i=1}^{m_t} \mu_i^{\mu_i r}. \end{aligned}$$

Let $\mu = \sum_{i=1}^{m_t} \mu_i$ be the sum of weights of the action at the fixed point component $X^{\zeta,t}$. Expanding out the product we obtain

$$(22) \quad \prod_{i=1}^{m_t} (\xi + c_1(\nu_{X^{\zeta,t},i})/\mu_i)^{\mu_i r} = \xi^{r\mu} + \xi^{r\mu-1} \left(r \sum_{i=1}^k c_1(\nu_{X^{\zeta,t},i}) \right) \\ + \xi^{r\mu-2} \left(r^2 \sum_{i \neq j} c_1(\nu_{X^{\zeta,t},i}) c_1(\nu_{X^{\zeta,t},j}) + \sum_{i=1}^k r(r-1/\mu_i) c_1(\nu_{X^{\zeta,t},i})^2 \right) + \dots$$

and $\dots$ indicates further terms with the property that the coefficient of $\xi^{r\mu-m}$ is polynomial in r . By the crepant assumption in Definition 4.1, the sum of the weights is $\mu = \sum_{i=1}^{m_t} \mu_i = 0$. Summing the terms from $\mathcal{S}^r, r \in \mathbb{Z}$ we obtain a distribution almost everywhere vanishing: For any singular value $t \in (-1, 1)$,

$$(23) \quad \sum_{r \in \mathbb{Z}} q^{d+\delta r} \tau_{X,\zeta,d+\delta r,t}(\alpha) = \sum_{r \in \mathbb{Z}} \prod_{i=1}^k \mu_i^{\mu_i r} q^{d+r\delta} \int_{[\mathcal{M}_n^{G,\zeta}(C,X,L_t,\zeta,d)]} \\ \left(1 + \xi^{-1} \sum_{i=1}^k r c_1(\nu_{X^{\zeta,t},i}) + \dots \right) \text{ev}^*(\alpha, \dots, \alpha)$$

where as before the terms $\dots$ are polynomial in the r . Now for any polynomial $f(r)$ in r ,

$$\sum_{r \in \mathbb{Z}} f(r) \left(\prod_{i=1}^k \mu_i^{\mu_i} q^\delta \right)^r =_{a.e.} 0$$

vanishes almost everywhere in q , being a function times a sum of derivatives of delta functions in q , see Remark 4.3. Since $\tau_{X/+G} - \tau_{X/-G}$ is a sum of such wall-crossing terms, this completes the proof of Theorem 1.12. $\square$

Remark 4.4. (a) The proof shows that if in fact the action of G on X is Calabi-Yau in the sense that $c_1^{\mathbb{C}^\times}(\nu_{X^{\zeta,t}}) = 0$, then the wall-crossing term is not only almost everywhere vanishing but also a sum of *at least second* derivatives of delta functions.

(b) The standard formulation of the crepant transformation conjecture in Coates-Ruan [13] etc. uses analytic continuation. The above results say nothing about convergence of the gauged Gromov-Witten potentials, so it is rather difficult to put the version above in this language. However, if the potentials have expressions as analytic functions with overlapping regions of definition on the torus $H_G^2(X, \mathbb{Q})/H_G^2(X, \mathbb{Z})$ with coordinate q , then they are equal on that region.

Example 4.5. (Crepant resolution of $\mathbb{C}^k/\mathbb{Z}_k$) Let $G = \mathbb{C}^\times$ act on $X = \mathbb{C}^{k+1}$ with weights $1, 1, 1, \dots, 1, -k$, and let $L_\pm$ denote the trivial line bundles with characters ± 1 . Then $X//_-G$ is isomorphic to the quotient stack $\mathbb{C}^k/\mathbb{Z}_k$ and $X//_+G$ is the crepant resolution of its coarse moduli space. Theorem 1.12 implies that the Gromov-Witten theories of $X//_\pm G$ are equivalent. Even in the case of an A_1 -singularity ($k = 2$) the

transformation relating the two equivalent Gromov-Witten theories is non-trivial; see Coates-Corti-Iritani-Tseng [14] and Bryan-Gholampour [9] for more on quantum cohomology of resolutions of ADE singularities.

Example 4.6. (Flop) Let $G = \mathbb{C}^\times$ act on $X = \mathbb{C}^{2k}$ with weights $1, -1$ each with multiplicity k , and let $L_\pm$ denote the trivial line bundles with characters ± 1 . Then $X//_t G$ undergoes a simple flop as t passes through zero (blow-down of a $\mathbb{P}^{k-1}$ followed by a blow-up) and Theorem 1.12 implies that the Gromov-Witten theories of $X//_\pm G$ are equivalent. See Iwao-Lee-Lin-Wang [32] for the invariance under simple flops in general and Example 1.14.

Example 4.7. (quantum McKay correspondences for abelian three-fold singularities) Let R be the regular representation of Γ and consider the affine scheme of representations of the McKay quiver with relations,

$$N = \{B \in \text{Hom}_\Gamma(R, \mathbb{C}^3 \otimes R) \mid B \wedge B = 0\}.$$

We denote by $\text{Irr}(\Gamma)$ the set of isomorphism classes of irreducible representations of Γ , for any $\rho \in \text{Irr}(\Gamma)$ let R_ρ denote the corresponding irreducible representation, and $R(\Gamma) = \bigoplus_{\rho \in \text{Irr}(\Gamma)} \mathbb{Z}\rho$ the representation ring. The group of equivariant automorphisms of the regular representation

$$\text{Aut}_\Gamma(R) = \prod_{\rho \in \text{Irr}(\Gamma)} GL(R_\rho)$$

acts on N by conjugation and induces a faithful action of $\mathbb{P} \text{Aut}_\Gamma(R) := \text{Aut}_\Gamma(R)/\mathbb{C}^\times$. Any *stability parameter* $\theta \in \text{Hom}_\mathbb{Z}(R(\Gamma), \mathbb{Z})$, $\theta(R) = 0$ induces a character

$$\chi_\theta : \mathbb{P} \text{Aut}_\Gamma(R) \rightarrow \mathbb{C}^\times, \quad [(g_\rho)_\rho] \mapsto \prod_{\rho \in \text{Irr}(\Gamma)} \det(g_\rho)^{\theta(\rho)}$$

and so a linearization of the trivial bundle. Denote

$$M_\theta := N//_\theta \mathbb{P} \text{Aut}_\Gamma(R)$$

the stack-theoretic quotient of the semistable locus in N with respect to θ . A result of Bridgeland, King, and Reid [6] shows that M_θ is a crepant resolution of the coarse moduli space of $\mathbb{C}^3/\Gamma$. Craw-Ishii [17] show further that any two projective crepant resolutions of (the coarse moduli space of) $\mathbb{C}^3/\Gamma$ are related by variation of git. By Theorem 1.12 and the results of Craw-Ishii [17], any two quotients $M_{\theta_-}, M_{\theta_+}$ for which stable=semistable have equivalent Gromov-Witten theories in the sense of Theorem 1.12 assuming that the git quotient at the wall being crossed is a smooth stack. This already follows from the results of Iwa-Lee-Lin-Wang [32], since as Craw-Ishii [17] shows, the transitions are all simple flops.

Remark 4.8. It would be interesting to understand, given a quiver with relations, for which stability parameters the git quotients are smooth stacks. For example, Ishii-Ueda [33] have given a non-degeneracy condition for smoothness of the moduli spaces associated to dimer models for generic stability parameters. For any codimension one wall whose associated git quotient is smooth (that is the semistable locus associated

to the polarization lying in a wall is smooth) one obtains an equivalence of Gromov-Witten theories. However, we do not know for which walls this condition holds.

REFERENCES

- [1] D. Abramovich, T. Graber, and A. Vistoli. Gromov-Witten theory of Deligne-Mumford stacks. *Amer. J. Math.*, 130(5):1337–1398, 2008.
- [2] D. Abramovich, T. Graber, M. Olsson, and H.-H. Tseng. On the global quotient structure of the space of twisted stable maps to a quotient stack. *J. Algebraic Geom.*, 16(4):731–751, 2007.
- [3] M. Ballard, D. Favero, and L. Katzarkov. Variation of geometric invariant theory quotients and derived categories. arXiv:1203.6643.
- [4] K. Behrend and B. Fantechi. The intrinsic normal cone. *Invent. Math.*, 128(1):45–88, 1997.
- [5] S. Boissière, É. Mann, and F. Perroni. Computing certain Gromov-Witten invariants of the crepant resolution of $\mathbb{P}(1, 3, 4, 4)$. *Nagoya Math. J.*, 201:1–22, 2011.
- [6] T. Bridgeland, A. King, and M. Reid. The McKay correspondence as an equivalence of derived categories. *J. Amer. Math. Soc.*, 14(3):535–554 (electronic), 2001.
- [7] M. Brion and C. Procesi. Action d’un tore dans une variété projective. In A. Connes et al., editors, *Operator Algebras, Unitary Representations, Enveloping Algebras, and Invariant Theory*, volume 62 of *Progress in Mathematics*, pages 509–539, Paris, 1989, 1990. Birkhäuser, Boston.
- [8] J. Bryan and A. Gholampour. Hurwitz-Hodge integrals, the E_6 and D_4 root systems, and the crepant resolution conjecture. *Adv. Math.*, 221(4):1047–1068, 2009.
- [9] J. Bryan and A. Gholampour. Root systems and the quantum cohomology of ADE resolutions. *Algebra Number Theory*, 2(4):369–390, 2008.
- [10] J. Bryan and T. Graber. The crepant resolution conjecture. In *Algebraic geometry—Seattle 2005. Part 1*, volume 80 of *Proc. Sympos. Pure Math.*, pages 23–42. Amer. Math. Soc., Providence, RI, 2009.
- [11] J. Bryan, T. Graber, and R. Pandharipande. The orbifold quantum cohomology of $\mathbb{C}^2/Z_3$ and Hurwitz-Hodge integrals. *J. Algebraic Geom.*, 17(1):1–28, 2008.
- [12] K. Cieliebak and D. Salamon. Wall crossing for symplectic vortices and quantum cohomology. *Math. Ann.*, 335(1):133–192, 2006.
- [13] T. Coates and Y. Ruan. Quantum cohomology and crepant resolutions: A conjecture. arXiv.org:0710.5901.
- [14] T. Coates, A. Corti, H. Iritani, and H. -H. Tseng. Computing genus-zero twisted Gromov-Witten invariants. *Duke Math. J.*, 147(3):377–438, 2009.
- [15] T. Coates, H. Iritani, and H. -H. Tseng. Wall-crossings in toric Gromov-Witten theory. I. Crepant examples. *Geom. Topol.*, 13(5):2675–2744, 2009.
- [16] B. Crauder and R. Miranda. Quantum cohomology of del Pezzo surfaces. preprint.
- [17] A. Craw and A. Ishii. Flops of G -Hilb and equivalences of derived categories by variation of GIT quotient. *Duke Math. J.*, 124(2):259–307, 2004.
- [18] J. de Jong and other authors. The Stacks Project. available at <http://math.columbia.edu/algebraic-geometry/stacks-git/>.
- [19] J.-M. Drezet and M. S. Narasimhan. Groupe de Picard des variétés de modules de fibrés semi-stables sur les courbes algébriques. *Invent. Math.*, 97(1):53–94, 1989.
- [20] I. V. Dolgachev and Y. Hu. Variation of geometric invariant theory quotients. *Inst. Hautes Études Sci. Publ. Math.*, (87):5–56, 1998. With an appendix by Nicolas Ressayre.
- [21] D. Edidin. Riemann-Roch for Deligne-Mumford stacks. 1205.4742.
- [22] G. Ellingsrud and L. Göttsche. Variation of moduli spaces and Donaldson invariants under change of polarization. *J. Reine Angew. Math.*, 467:1–49, 1995.
- [23] A. R. P. Gaio and D. A. Salamon. Gromov-Witten invariants of symplectic quotients and adiabatic limits. *J. Symplectic Geom.*, 3(1):55–159, 2005.

- [24] A. B. Givental. Equivariant Gromov-Witten invariants. *Internat. Math. Res. Notices*, (13):613–663, 1996.
- [25] E. Gonzalez and C. Woodward. Quantum Witten localization. arXiv:0811.3358.
- [26] E. Gonzalez and C. Woodward. Quantum cohomology of toric orbifolds. arXiv:1207.3253.
- [27] T. Graber and R. Pandharipande. Localization of virtual classes. *Invent. Math.*, 135(2):487–518, 1999.
- [28] V. Guillemin and S. Sternberg. Birational equivalence in the symplectic category. *Invent. Math.*, 97(3):485–522, 1989.
- [29] D. Halpern-Leistner. The derived category of a GIT quotient. arXiv:1203.0276.
- [30] L. Hörmander. *The Analysis of Linear Partial Differential Operators I*, volume 256 of *Grundlehren der mathematischen Wissenschaften*. Springer-Verlag, Berlin-Heidelberg-New York, second edition, 1990.
- [31] Y. Hu and S. Keel. Mori dream spaces and GIT. *Michigan Math. J.*, 48:331–348, 2000. Dedicated to William Fulton on the occasion of his 60th birthday.
- [32] Y. Iwao, Y.-P. Lee, H.-W. Lin, and C.-L. Wang. Invariance of Gromov-Witten theory under a simple flop. *J. Reine Angew. Math.*, 663:67–90, 2012.
- [33] A. Ishii and K. Ueda. Dimer models and exceptional collections. arXiv:0911.4529.
- [34] J. Kalkman. Cohomology rings of symplectic quotients. *J. Reine Angew. Math.*, 485:37–52, 1995.
- [35] Y. Kawamata. D -equivalence and K -equivalence. *J. Differential Geom.*, 61(1):147–171, 2002.
- [36] G. Kempf and L. Ness. The length of vectors in representation spaces. In K. Lønsted, editor, *Algebraic Geometry*, volume 732 of *Lecture Notes in Mathematics*, pages 233–244, Copenhagen, 1978, 1979. Springer-Verlag, Berlin-Heidelberg-New York.
- [37] Y. -H. Kiem and J. Li. Virtual wall crossing formulas for C^* -flips, 2009. arXiv:0905.4770.
- [38] F. C. Kirwan. *Cohomology of Quotients in Symplectic and Algebraic Geometry*, volume 31 of *Mathematical Notes*. Princeton Univ. Press, Princeton, 1984.
- [39] M. Kontsevich. Enumeration of rational curves via torus actions. In *The moduli space of curves (Texel Island, 1994)*, pages 335–368. Birkhäuser Boston, Boston, MA, 1995.
- [40] Y. -P. Lee, H. -W. Lin, and C. -L. Wang. Flops, motives, and invariance of quantum rings. *Ann. of Math. (2)*, 172(1):243–290, 2010.
- [41] E. Lerman. Symplectic cuts. *Math. Res. Letters*, 2:247–258, 1995.
- [42] B. H. Lian, K. Liu, and S.-T. Yau. Mirror principle. I. *Asian J. Math.*, 1(4):729–763, 1997.
- [43] A. -M. Li and Y. Ruan. Symplectic surgery and Gromov-Witten invariants of Calabi-Yau 3-folds. *Invent. Math.*, 145(1):151–218, 2001.
- [44] Yu. I. Manin. *Cubic forms*, volume 4 of *North-Holland Mathematical Library*. North-Holland Publishing Co., Amsterdam, second edition, 1986. Algebra, geometry, arithmetic, Translated from the Russian by M. Hazewinkel.
- [45] J. McKernan. Mori dream spaces. *Jpn. J. Math.*, 5(1):127–151, 2010.
- [46] D. Mumford, J. Fogarty, and F. Kirwan. *Geometric Invariant Theory*, volume 34 of *Ergebnisse der Mathematik und ihrer Grenzgebiete, 2. Folge*. Springer-Verlag, Berlin-Heidelberg-New York, third edition, 1994.
- [47] I. Mundet i Riera. A Hitchin-Kobayashi correspondence for Kähler fibrations. *J. Reine Angew. Math.*, 528:41–80, 2000.
- [48] M. Popa and M. Roth. Stable maps and Quot schemes. *Invent. Math.*, 152(3):625–663, 2003.
- [49] M. Reid. Minimal models of canonical 3-folds. In *Algebraic varieties and analytic varieties (Tokyo, 1981)*, volume 1 of *Adv. Stud. Pure Math.*, pages 131–180. North-Holland, Amsterdam, 1983.
- [50] A. H. W. Schmitt. *Geometric invariant theory and decorated principal bundles*. Zurich Lectures in Advanced Mathematics. European Mathematical Society (EMS), Zürich, 2008.

- [51] E. Segal. Equivalences Between GIT Quotients of Landau-Ginzburg B-Models. *Communications in Mathematical Physics, Volume 304, Issue 2, pp.411-432*, 304:411–432, June 2011. arxiv:0910.5534.
- [52] M. Thaddeus. Geometric invariant theory and flips. *J. Amer. Math. Soc.*, 9(3):691–723, 1996.
- [53] B. Toen. Théorèmes de Riemann-Roch pour les champs de Deligne-Mumford. *K-Theory*, 18(1):33–76, 1999.
- [54] E. Witten. Phases of $N = 2$ theories in two dimensions. *Nuclear Phys. B*, 403(1-2):159–222, 1993.
- [55] C. Woodward. Quantum Kirwan morphism and Gromov-Witten invariants of quotients. arXiv:1204.1765.

DEPARTMENT OF MATHEMATICS UNIVERSITY OF MASSACHUSETTS BOSTON 100 WILLIAM T. MORRISSEY BOULEVARD BOSTON, MA 02125

E-mail address: `eduardo@math.umb.edu`

MATHEMATICS-HILL CENTER, RUTGERS UNIVERSITY, 110 FRELINGHUYSEN ROAD, PISCATAWAY, NJ 08854-8019, U.S.A.

E-mail address: `ctw@math.rutgers.edu`